

FACTORIZATION OF HOMOTOPIES OF NANOPHRASES

ANDREW GIBSON

ABSTRACT. Homotopy on nanophrases is an equivalence relation defined using some data called a homotopy data triple. We define a product on homotopy data triples. We show that any homotopy data triple can be factorized into a product of prime homotopy data triples and this factorization is unique up to isomorphism and order. If a homotopy data triple is composite, we show that equivalence of nanophrases under the corresponding homotopy can be calculated just by using the homotopies given by its prime factors.

1. INTRODUCTION

A word is a sequence of letters. If every letter that appears in a word appears exactly twice, then the word is a Gauss word. Let α be a fixed set. A nanoword over α is a Gauss word paired with a map from the set of letters appearing in this word to α . A Gauss phrase is a sequence of words such that their concatenation forms a Gauss word. A nanophrase over α is a Gauss phrase paired with a map from the set of letters appearing in this phrase to α . Turaev defined nanowords in [15] and nanophrases in [14].

For nanowords and nanophrases, Turaev defined moves which are determined by an involution on α called τ and a subset of $\alpha \times \alpha \times \alpha$ called S . We call the triple (α, τ, S) a homotopy data triple. Fixing a homotopy data triple, the moves generate an equivalence relation on nanowords and nanophrases over α called homotopy. Different homotopy data triples can give different equivalence relations.

Two homotopy data triples, (α, τ, S) and (α', τ', S') are isomorphic if there is a bijection from α to α' which transforms τ into τ' and S into S' . Isomorphic homotopy data triples give equivalent homotopies [14].

We define a product on homotopy data triples. A homotopy data triple is *composite* if it can be represented as a non-trivial product of homotopy data triples. If not, the homotopy data triple is *prime*. We show that any homotopy data triple can be represented uniquely, up to order and isomorphism, as a product of prime homotopy data triples (Proposition 4.7).

In this paper, our main aim is to show that for any composite homotopy data triple, the study of the homotopy it gives can be reduced to the study of the homotopies given by its prime factors.

Let (α, τ, S) be a composite homotopy data triple. Let $\mathcal{P}_R(\alpha)$ be the set of nanophrases over α satisfying the following conditions: (1) each component is associated with a prime factor of (α, τ, S) and any letters in that component map to the factor; and (2) adjacent components are associated with different prime factors. Any nanoword over α can be uniquely split into a nanophrase in $\mathcal{P}_R(\alpha)$ such that each component is non-empty. This gives a map from nanowords over α to $\mathcal{P}_R(\alpha)$ called the decomposing map.

Date: January 30, 2010.

2000 Mathematics Subject Classification. Primary 57M99; Secondary 68R15.

Key words and phrases. nanowords, nanophrases, homotopy invariant.

This work was supported by a Scholarship from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

We define two equivalence relations on $\mathcal{P}_R(\alpha)$. The equivalence relation $\sim_K$ is a restriction of homotopy to $\mathcal{P}_R(\alpha)$. It can be defined entirely in terms of the homotopies associated with the prime factors of (α, τ, S) . The equivalence relation $\sim_P$ is generated by $\sim_K$, reductions (moves on nanophrases which allow us to remove empty components) and their inverses.

The key result of this paper is that there is a bijection between the set of homotopy classes of nanowords over α and the set of equivalence classes of $\mathcal{P}_R(\alpha)$ under $\sim_P$ (Theorem 5.8).

We use this bijection to define a homotopy invariant of nanowords. Given a nanoword over α , we apply the decomposing map to get a nanophrase p in $\mathcal{P}_R(\alpha)$. We say that p is reducible if it is equivalent under $\sim_K$ to a nanophrase q with an empty component. By applying a reduction to q we get a nanophrase which is equivalent to p under $\sim_P$ but has less components. By repeating this process we will eventually get an irreducible nanophrase. We show that, irrespective of how we make the reductions, the irreducible nanophrase that we obtain is unique up to equivalence under $\sim_K$ (Proposition 5.14). In fact, this gives a complete homotopy invariant of the original nanoword (Theorem 5.15). We generalize these results to nanophrases in Section 8.

In general, calculating this invariant can be hard, because it is not always easy to tell whether or not a nanophrase is irreducible. Moreover, using the invariant to show that different nanowords are not homotopic entails showing that their irreducible nanophrases are not equivalent under $\sim_K$ which, in general, is also difficult. However, the calculation of the invariant of a nanoword and determination of the equivalence of this invariant for different nanowords can be achieved just by using the homotopies given by the prime factors of (α, τ, S) .

A homotopy of nanophrases over α is *equality decidable* if there exists a finite time algorithm which given any two nanophrases over α , determines whether or not they are homotopic. The homotopy is *reduction decidable* if there exists a finite time algorithm which given any nanophrase over α , determines whether or not it is reducible.

In the case where S is the empty set, for any α and τ , the homotopy given by (α, τ, S) is reduction and equality decidable (Proposition 6.6). In all other cases, determining whether a homotopy is reduction or equality decidable is an open problem. On the other hand, we use our complete invariant of nanowords and nanophrases to show the following fact. Let (α, τ, S) be a composite homotopy data triple and let its prime factors be denoted by (α_i, τ_i, S_i) . If the homotopies given by each (α_i, τ_i, S_i) are all reduction and equality decidable, the homotopy given by (α, τ, S) is reduction and equality decidable (Theorem 8.7).

In Section 7 we examine several invariants which can be used to give sufficient conditions for irreducibility. Two of the invariants we consider, U and V , are new. Both invariants are generalizations of our S_o invariant for Gauss phrases [5]. The V invariant is defined for all homotopies but U is just defined for homotopies where S is diagonal (that is, S is of the form $\{(a, a, a) \mid a \in \alpha\}$). The U invariant is a generalization to nanophrases of Turaev's self-linking function for nanowords [15]. Having written this paper we discovered that Fukunaga had independently generalized our S_o invariant [3]. We give a proof that Fukunaga's invariant and our U invariant are equivalent in Section 7.6.

Acknowledgments. The author is very grateful for all the help, advice and encouragement given to him by his supervisor Hitoshi Murakami. He would like to thank Tomonori Fukunaga for several useful conversations. The author is also very grateful for advice he received from Vladimir Turaev.

2. NANOWORDS AND NANOPHRASES

Nanowords and nanophrases were defined by Turaev in [15] and [14]. In this section we briefly recall various definitions from those papers.

An *alphabet* is a finite set and its elements are called *letters*. For any positive integer m , let $\hat{m}$ denote the set $\{1, 2, \dots, m\}$. A *word* on an alphabet $\mathcal{A}$ of length m is a map from $\hat{m}$ to $\mathcal{A}$. Informally, we can think of a word w on $\mathcal{A}$ as a finite sequence of letters in $\mathcal{A}$ and we will usually write words in this way. So, for example, $ABCA$ is a word of length 4 on $\{A, B, C, D\}$. We can write this word as a map from $\hat{4}$ to $\{A, B, C, D\}$, where 1 maps to A , 2 maps to B , 3 maps to C and 4 maps to A . The empty word of length 0 on any alphabet is written $\emptyset$.

An n -component *phrase* on an alphabet $\mathcal{A}$ is a sequence of n words on $\mathcal{A}$. We write phrases as a sequence of words, separated by ‘|’ symbols. For example, $A|BCA|CD$ is a 3-component phrase on the alphabet $\{A, B, C, D\}$. Note that in [14], Turaev always encloses phrases in brackets but in this paper we omit them.

A *Gauss word* on an alphabet $\mathcal{A}$ is a word on $\mathcal{A}$ such that every letter in $\mathcal{A}$ appears exactly twice in the word. A *Gauss phrase* on an alphabet $\mathcal{A}$ is a phrase on $\mathcal{A}$ such that every letter in $\mathcal{A}$ appears exactly twice in the phrase. Equivalently, a Gauss phrase on $\mathcal{A}$ is a phrase on $\mathcal{A}$ such that the concatenation of the words appearing in the phrase is a Gauss word on $\mathcal{A}$. By definition, the single component appearing in a 1-component Gauss phrase is a Gauss word.

Let α be a non-empty set. An α -alphabet is an alphabet $\mathcal{A}$ which has an associated map from $\mathcal{A}$ to α . This map is called a *projection*. Following Turaev, we will often write $|X|$ for the image of X under the projection, where X is a letter in $\mathcal{A}$. An *isomorphism* of α -alphabets $\mathcal{A}_1$ and $\mathcal{A}_2$ is a bijection f from $\mathcal{A}_1$ to $\mathcal{A}_2$ such that $|f(X)|$ is equal to $|X|$ for all letters X in $\mathcal{A}_1$.

A nanoword over α is a pair $(\mathcal{A}, w)$ where $\mathcal{A}$ is an α -alphabet and w is a Gauss word on $\mathcal{A}$. We write $\mathcal{N}(\alpha)$ for the set of nanowords over α .

An n -component *nanophrase over α* is a pair $(\mathcal{A}, p)$ where $\mathcal{A}$ is an α -alphabet and p is an n -component Gauss phrase on $\mathcal{A}$. There is a unique nanophrase with 0 components which we write $\emptyset_P$. Note that in [14] Turaev uses $\emptyset$ to denote an 0-component nanophrase and $(\emptyset)$ to represent a nanophrase with a single empty component. We always use $\emptyset$ to represent an empty word and therefore, according to context, it also represents an empty Gauss word or a nanophrase with a single empty component.

We write $\mathcal{P}_n(\alpha)$ for the set of n -component nanophrases over α and write $\mathcal{P}(\alpha)$ for the set of all nanophrases over α . Thus, $\mathcal{P}(\alpha)$ is the union of the $\mathcal{P}_n(\alpha)$ with n running from 0 to infinity.

There is a natural bijection between nanowords over α and 1-component nanophrases over α induced by mapping a Gauss word w to a 1-component Gauss phrase containing a single component w . In this paper, we will consider nanowords over α and 1-component nanophrases over α to be the same. Thus $\mathcal{N}(\alpha)$ and $\mathcal{P}_1(\alpha)$ represent the same set.

Two nanophrases over α , $(\mathcal{A}_1, p_1)$ and $(\mathcal{A}_2, p_2)$, are *isomorphic* if there exists an isomorphism of α -alphabets f from $\mathcal{A}_1$ to $\mathcal{A}_2$ such that f applied letterwise to the i th component of p_1 gives the i th component of p_2 for all i .

Let τ be an involution on α (that is, $\tau(\tau(a))$ is equal to a for all a in α). Let S be a subset of $\alpha \times \alpha \times \alpha$. We say that S is *diagonal* if $S = \{(a, a, a) \mid a \in \alpha\}$.

Turaev defined *homotopy moves* for nanophrases on α . In moves on nanowords, the lower case letters x, y, z and t represent arbitrary sequences of letters, possibly including one or more ‘|’ symbols, so that the phrases on each side of the move are Gauss phrases. The moves are

move H1: for any $|A|$,

$$\begin{aligned}
& (\mathcal{A}, xAAy) \longleftrightarrow (\mathcal{A} - \{A\}, xy) \\
\text{move H2: if } \tau(|A|) = |B|, & \\
& (\mathcal{A}, xABYBAz) \longleftrightarrow (\mathcal{A} - \{A, B\}, xyz) \\
\text{move H3: if } (|A|, |B|, |C|) \in S, & \\
& (\mathcal{A}, xABYACzBCt) \longleftrightarrow (\mathcal{A}, xBAyCAzCBt)
\end{aligned}$$

where the projections of the α -alphabets on the right hand side of the moves H1 and H2 are restrictions of the corresponding α -alphabets on the left hand side.

Two nanophrases are *homotopic* if there exists a finite sequence of homotopy moves and isomorphisms which transform one into the other. As none of the moves add or remove components, the number of components of a nanophrase is a homotopy invariant.

Note that if we change α , τ or S we may get a different kind of homotopy. We call the triple (α, τ, S) a *homotopy data triple* (note that Turaev uses the term *homotopy data* [15]).

Two homotopy data triples, (α, τ, S) and (α', τ', S') are *isomorphic* if there exists a bijection f from α to α' such that $\tau' \circ f$ is equivalent to $f \circ \tau$ and (a, b, c) is in S if and only if $(f(a), f(b), f(c))$ is in S' . Such an isomorphism induces a map from $\mathcal{P}(\alpha)$ to $\mathcal{P}(\alpha')$ which is homotopy preserving [14]. Thus the homotopies given by isomorphic homotopy data triples are equivalent.

Let p be a nanophrase. We write $\text{nc}(p)$ for the number of components in p and define $c_i(p)$ to be the i th component of p . The *rank* of p is the number of distinct letters appearing in p . We write it $\text{rank}(p)$. The *homotopy rank* of p , which is written $\text{hr}(p)$, is the minimum rank attained by a nanophrase in the homotopy class of p . We say that a nanophrase p is *minimal* if $\text{rank}(p)$ is equal to $\text{hr}(p)$.

A nanoword is *contractible* if it is homotopic to the empty word $\emptyset$. Thus, for a contractible nanoword w , $\text{hr}(w)$ is 0.

3. SOME LEMMAS ON NANOPHRASES

We define a map χ from $\mathcal{P}(\alpha)$ to $\mathcal{N}(\alpha)$ as follows. For a nanophrase p in $\mathcal{P}(\alpha)$ we define $\chi(p)$ to be the nanoword obtained by concatenating the components of p . That is $w_1|w_2|\dots|w_n$ is mapped to $w_1w_2\dots w_n$. By definition $\chi(\emptyset_P)$ is $\emptyset$. We call χ the *concatenating map*. The following lemma is Lemma 4.3 of [4].

Lemma 3.1 (Fukunaga). *Let p_1 and p_2 be nanophrases in $\mathcal{P}(\alpha)$. If $p_1 \sim p_2$, then $\chi(p_1) \sim \chi(p_2)$.*

Note that in [4] the lemma above was stated in the case where S is diagonal. However, it trivially extends to the general case.

Let O be a subset of $\hat{n}$. Then we define $\mathcal{P}_n(\alpha, O)$ to be the subset of $\mathcal{P}_n(\alpha)$ where p is in $\mathcal{P}_n(\alpha, O)$ if $c_i(p)$ is $\emptyset$ for all i in O . Let $\sim_O$ be the equivalence relation generated by isomorphisms and by homotopy moves which relate elements in $\mathcal{P}_n(\alpha, O)$. Clearly $p \sim_O p'$ implies $p \sim p'$.

We define a map f_O from $\mathcal{P}_n(\alpha)$ to $\mathcal{P}_n(\alpha, O)$ by saying that $f_O(p)$ is the nanophrase derived from p by deleting all letters that appear at least once in any component with index in O .

Lemma 3.2. *Let p and p' be two nanophrases in $\mathcal{P}_n(\alpha)$. If $p \sim p'$ then $f_O(p) \sim_O f_O(p')$.*

Proof. It is enough to check that if p and p' are related by a single isomorphism or homotopy move then $f_O(p) \sim_O f_O(p')$.

If p and p' are isomorphic then $f_O(p)$ and $f_O(p')$ are also isomorphic.

Suppose that p and p' are related by an H1 move. We assume that the move removes a letter A from p . If A appeared in a component with index in O , then A

will not appear in $f_O(p)$ and so $f_O(p)$ is $f_O(p')$. Otherwise, $f_O(p)$ and $f_O(p')$ are related by an H1 move.

Now suppose that p and p' are related by an H2 move. We assume that the move removes letters A and B from p . If A appeared in a component with index in O , then B also appears in such a component and so A and B will both be deleted in $f_O(p)$. Thus $f_O(p)$ is $f_O(p')$. If A does not appear in a component with index in O , then neither does B and $f_O(p)$ and $f_O(p')$ are related by an H2 move.

Finally suppose that p and p' are related by an H3 move. The move affects three letters. Let k be the number of letters affected by the move which appear in a component with index in O . Note that if one of the letters affected by the move appears in a component with index in O , then at least one of the other two letters must also appear in the same component. Thus k must be 0, 2 or 3. If k is 2 or 3 then $f_O(p)$ is $f_O(p')$. If k is 0 then $f_O(p)$ and $f_O(p')$ are related by an H3 move. $\square$

Lemma 3.3. *Let p and p' be two nanophrases in $\mathcal{P}_n(\alpha, O)$. If $p \sim p'$ then $p \sim_O p'$.*

Proof. If $p \sim p'$ then there exists a sequence of nanophrases in $\mathcal{P}_n(\alpha)$, where each consecutive pair in the sequence is related by an isomorphism or a single homotopy move. By applying f_O to each nanophrase in the sequence, we get a sequence of nanophrases in $\mathcal{P}_n(\alpha, O)$ showing $p \sim_O p'$. $\square$

Lemma 3.4. *Let p and p' be two nanophrases in $\mathcal{P}_n(\alpha)$ and let O be a subset of the set of indices of empty components in p' . If $p \sim p'$, then $p \sim f_O(p)$.*

Proof. We have

$$f_O(p) \sim_O f_O(p') = p' \sim p,$$

where the first relation is given by Lemma 3.2, the second relation follows from the definition of f_O and the third relation is the assumption. $\square$

Lemma 3.5. *Let p be a nanophrase in $\mathcal{P}_n(\alpha)$ such that $\text{rank}(p) = \text{hr}(p)$. Let p' be a nanophrase in $\mathcal{P}_n(\alpha)$ such that the i th component of p' is empty. If $p \sim p'$, then the i th component of p is empty.*

Proof. Suppose the i th component of p is not empty. Let O be the set $\{i\}$. Then by Lemma 3.4, $p \sim f_O(p)$. Now $\text{rank}(f_O(p))$ is less than $\text{rank}(p)$, as the i th component of p is not empty. Then we have

$$\text{hr}(p) \leq \text{rank}(f_O(p)) < \text{rank}(p) = \text{hr}(p),$$

which is a contradiction. Thus the i th component of p must be empty. $\square$

We write $O(p)$ for the set of indices of components that are empty in p .

Lemma 3.6. *Let p and p' be two nanophrases in $\mathcal{P}_n(\alpha)$. If $p \sim p'$ and $\text{rank}(p) = \text{rank}(p') = \text{hr}(p)$ then $O(p) = O(p')$.*

Proof. For any element i in $O(p')$, Lemma 3.5 implies i is in $O(p)$. In other words $O(p') \subseteq O(p)$. By symmetry, we get the opposite inclusion and so $O(p) = O(p')$. $\square$

Let p be an n -component nanophrase and let O be a subset of $\hat{n}$ of size $|O|$. Define $x(p, O)$ to be the $n - |O|$ component nanophrase derived from p by deleting all components with index appearing in O and all letters which appear in those components.

Lemma 3.7. *Let p and p' be two nanophrases in $\mathcal{P}_n(\alpha)$. If $p \sim p'$ then $x(p, O) \sim x(p', O)$.*

Proof. Let p_1 and p_2 be two nanophrases in $\mathcal{P}_n(\alpha, O)$. If $p_1 \sim_O p_2$ then $x(p_1, O) \sim x(p_2, O)$. The result then follows from Lemma 3.2. $\square$

Lemma 3.7 corresponds to the well-known fact that a sub-link is an invariant of a link.

4. FACTORIZATIONS OF HOMOTOPY DATA TRIPLES

Let (α, τ, S) be a homotopy data triple. From now on, we will assume that α is finite. Let β be a τ -invariant subset of α and γ be the set $\alpha - \beta$. Note that γ is also τ -invariant. We define τ_β to be the restriction of τ to β and τ_γ to be the restriction of τ to γ . We define S_β to be the intersection of S and $\beta \times \beta \times \beta$ and S_γ to be the intersection of S and $\gamma \times \gamma \times \gamma$. If every element of S appears either in S_β or S_γ (that is, S is equal to $S_\beta \cup S_\gamma$), we say that $(\beta, \tau_\beta, S_\beta)$ is a *factor* of (α, τ, S) . Note that if $(\beta, \tau_\beta, S_\beta)$ is a factor of (α, τ, S) , so is $(\gamma, \tau_\gamma, S_\gamma)$.

Any homotopy data triple (α, τ, S) is a factor of itself, as α is a τ -invariant subset of α and S is contained in $\alpha \times \alpha \times \alpha$. The homotopy data triple $(\emptyset, \tau_\emptyset, \emptyset)$, where $\tau_\emptyset$ is the empty map, is also a factor of any homotopy data triple. We note however that $(\emptyset, \tau_\emptyset, \emptyset)$ does not give a homotopy because the concept of an α -alphabet is not defined when α is $\emptyset$.

A factor $(\beta, \tau_\beta, S_\beta)$ of a homotopy data triple (α, τ, S) is said to be a *proper factor* of (α, τ, S) if β is a non-empty proper subset of α . We say that a homotopy data triple (α, τ, S) is *composite* if it has a proper factor. If a homotopy data triple (α, τ, S) has no proper factors and α is non-empty, then we say that the homotopy data triple is *prime*.

Example 4.1. Let α be the set $\{a, b, c, d\}$. Let τ be the map which swaps a with b and c with d . Let S be the set $\{(a, b, a), (c, d, c)\}$. Now α has two orbits under τ , so if (α, τ, S) has a proper factor $(\beta, \tau_\beta, S_\beta)$, then β must contain one of those orbits. Without loss of generality, we assume β is the set $\{a, b\}$. Set γ to be the set $\{c, d\}$. Then τ_β , the restriction of τ to β , is the map which swaps a with b . The restriction of τ to γ , τ_γ , is the map that swaps c with d . The set S_β , the intersection of S and $\beta \times \beta \times \beta$, is $\{(a, b, a)\}$. The set S_γ , the intersection of S and $\gamma \times \gamma \times \gamma$, is $\{(c, d, c)\}$. Now S is equal to $S_\beta \cup S_\gamma$, so $(\beta, \tau_\beta, S_\beta)$ is a proper factor of (α, τ, S) . Thus (α, τ, S) is composite.

Example 4.2. Let α and τ be the same as in Example 4.1 and let S be the set $\{(a, b, c), (b, c, d)\}$. As before, α has two orbits under τ , so if (α, τ, S) has a proper factor $(\beta, \tau_\beta, S_\beta)$, then β must contain one of those orbits. We assume β is the set $\{a, b\}$. Then γ , τ_β and τ_γ are the same as in Example 4.1. However, in this case both S_β and S_γ are empty. Clearly S is not equal to $S_\beta \cup S_\gamma$, so $(\beta, \tau_\beta, S_\beta)$ is not a factor of (α, τ, S) . Thus (α, τ, S) is prime.

We remark that if α has only one orbit under τ then the homotopy data triple (α, τ, S) must be prime.

In the following example we consider the case where S is diagonal.

Example 4.3. Let α_G be the set $\{a\}$, let τ_G be the identity map and let S_G be the set $\{(a, a, a)\}$. Let α_F be the set $\{a, b\}$, let τ_F be the map swapping a and b , and let S_F be the set $\{(a, a, a), (b, b, b)\}$. Then both (α_G, τ_G, S_G) and (α_F, τ_F, S_F) are prime. We note that (α_G, τ_G, S_G) gives the open Gauss phrase homotopy and the homotopy data triple (α_F, τ_F, S_F) gives the open flat virtual link homotopy (see, for example, [5] or [14]).

Consider the general case of a homotopy data triple (α, τ, S) where S is diagonal. It is easy to check that if (α, τ, S) is prime, then it is isomorphic to (α_G, τ_G, S_G) or (α_F, τ_F, S_F) . Otherwise, (α, τ, S) is composite and has a proper factor which is isomorphic to either (α_G, τ_G, S_G) or (α_F, τ_F, S_F) .

We now define the product of two homotopy data triples. Let (α_1, τ_1, S_1) and (α_2, τ_2, S_2) be homotopy data triples. Let α be the subset of $\{1, 2\} \times (\alpha_1 \cup \alpha_2)$ given by

$$\{(1, a) \mid a \in \alpha_1\} \cup \{(2, a) \mid a \in \alpha_2\}.$$

We define τ to be the involution on α where (i, a) maps to $(i, \tau_i(a))$ for each (i, a) in α . We define S to be the subset of $\alpha \times \alpha \times \alpha$ given by

$$\{((1, a), (1, b), (1, c)) \mid (a, b, c) \in S_1\} \cup \{((2, a), (2, b), (2, c)) \mid (a, b, c) \in S_2\}.$$

Then the product of (α_1, τ_1, S_1) and (α_2, τ_2, S_2) is (α, τ, S) .

We note that if α_1 and α_2 are disjoint, then the product of (α_1, τ_1, S_1) and (α_2, τ_2, S_2) is isomorphic to $(\alpha_1 \cup \alpha_2, \tau, S_1 \cup S_2)$ where $\tau(a)$ is defined to be $\tau_1(a)$ if a is in α_1 and $\tau_2(a)$ if a is in α_2 .

The product of homotopy data triples induces a product on the isomorphism classes of homotopy data triples. This product is well-defined because, if (α_1, τ_1, S_1) is isomorphic to $(\alpha'_1, \tau'_1, S'_1)$ and (α_2, τ_2, S_2) is isomorphic to $(\alpha'_2, \tau'_2, S'_2)$, then the product of (α_1, τ_1, S_1) and (α_2, τ_2, S_2) is isomorphic to the product of $(\alpha'_1, \tau'_1, S'_1)$ and $(\alpha'_2, \tau'_2, S'_2)$.

Suppose $(\beta, \tau_\beta, S_\beta)$ is a factor of (α, τ, S) . Let γ be the set $\alpha - \beta$, let τ_γ be the restriction of τ to γ and let S_γ be the intersection of S and $\gamma \times \gamma \times \gamma$. Then the product of $(\beta, \tau_\beta, S_\beta)$ and $(\gamma, \tau_\gamma, S_\gamma)$ is isomorphic to (α, τ, S) .

It is easy to check that up to isomorphism, the product on homotopy data triples is commutative and associative. The unit of the product is the triple $(\emptyset, \tau_\emptyset, \emptyset)$.

We now show that, up to isomorphism and reordering, any homotopy data triple can be uniquely written as a product of prime homotopy data triples. We first state some lemmas.

Lemma 4.4. *Suppose h is a homotopy data triple that can be represented as the product of f and g . Suppose k is a prime factor of h . Then k is a prime factor of f or g .*

Proof. In this proof, the data for a homotopy data triple x is written α_x , τ_x and S_x . As h can be represented as the product fg , α_f and α_g are τ_h -invariant sets which partition α_h .

Now suppose that $\alpha_k \cap \alpha_f$ and $\alpha_k \cap \alpha_g$ are both non-empty. Write α_l for $\alpha_k \cap \alpha_f$ and α_m for $\alpha_k \cap \alpha_g$. Then both α_l and α_m are τ_h -invariant sets. We define S_l as the intersection of S_h and $\alpha_l \times \alpha_l \times \alpha_l$ and define S_m as the intersection of S_h and $\alpha_m \times \alpha_m \times \alpha_m$. Since S_h is equal to $S_f \cup S_g$, S_k must equal $S_l \cup S_m$. This implies that (α_l, τ_l, S_l) is a factor of k where τ_l is the restriction of τ_h to α_l . However, this contradicts the primality of k . Thus α_k is wholly contained in either α_f or α_g .

Without loss of generality, we assume that α_k is a subset of α_f . Let α_p be the set $\alpha_f - \alpha_k$ and let S_p be the intersection of S_f and $\alpha_p \times \alpha_p \times \alpha_p$. Let α_q be the set $\alpha_h - \alpha_k$ and let S_q be the intersection of S_h and $\alpha_q \times \alpha_q \times \alpha_q$. Now suppose that $S_k \cup S_p$ is not equal to S_f . Then this would imply that $S_k \cup S_q$ is not equal to S_h . However, this contradicts the fact that k is a factor of h . Thus $S_k \cup S_p$ is equal to S_f and k is a factor of f . $\square$

Lemma 4.5. *Let g, h be homotopy data triples and k be a prime homotopy data triple. If the product of g and k is isomorphic to the product of h and k , then g and h are isomorphic.*

Proof. We write gk for the product of g and k and hk for the product of h and k . Let f be the isomorphism from gk to hk . As k is a prime factor of gk , the image of k , $f(k)$, is a prime factor of hk . Thus by Lemma 4.4, $f(k)$ is either a factor of h or of k .

If $f(k)$ is a factor of k , then $f(k)$ is k as they have the same size. Thus the image of g must be h and f restricted to g gives an isomorphism from g to h .

If $f(k)$ is a factor of h , then h can be written kh' for some h' . Then f maps g onto $h'k$ and this gives an isomorphism from g to $h'k$. Since the product of homotopy data triples is commutative up to isomorphism, $h'k$ is isomorphic to kh' . Thus g is isomorphic to h . $\square$

Lemma 4.6. *Let h be a prime homotopy data triple and h_i be a set of prime homotopy data triples, for i running from 1 to n . Let k be the product of the h_i . If h is a factor of k , then h is isomorphic to one of the h_i .*

Proof. We prove by induction on n . If n is 1 then k is isomorphic to h_1 which is prime. Therefore h must be isomorphic to k .

We now assume the statement is true for $n - 1$ and prove it for n . If h is isomorphic to h_1 then the statement is true. If not, Lemma 4.4 implies that h must be a factor of the product of h_i for i running from 2 to n . Then this product has $n - 1$ triples and so by assumption, h is isomorphic to one of the h_i for i not equal to 1. $\square$

We then have the following proposition.

Proposition 4.7. *Any homotopy data triple can be represented as a product of prime homotopy data triples. This representation is unique up to isomorphism and the order of the triples in the product.*

Proof. Existence of such a representation is easy to show. If the homotopy data triple is prime we already have our representation. If not, it is composite and can be written as the product of two homotopy data triples. We consider each of these in turn and factor any that are composite. We continue this process until only prime factors remain.

Let h be a homotopy data triple and suppose it can be factored into triples h_i for i running from 1 to m and also into triples h'_j for j running from 1 to n . Without loss of generality we may assume that n is greater than or equal to m .

We prove uniqueness by induction on m . For if m is 1 then h_1 is prime and is isomorphic to h . Thus n must also be 1 and h'_1 is isomorphic to h_1 .

We now assume the statement is true for $m - 1$ and prove it for m . By Lemma 4.6, h_1 must be isomorphic to one of the h'_j . By reordering the h'_j if necessary, we may assume that it is h'_1 . Then by Lemma 4.5 the product of h_i with i running from 2 to m and the product of h'_j for j running from 2 to n are isomorphic. As the product of h_i with i running from 2 to m has $m - 1$ factors, we use the induction assumption to show that n is equal to m and, after appropriate reordering of the h'_j , h_j is isomorphic to h'_j for each j . $\square$

Example 4.8. Let (α, τ, S) be a homotopy data triple with diagonal S . It is easy to check that (α, τ, S) is isomorphic to the product

$$(\alpha_G, \tau_G, S_G)^k (\alpha_F, \tau_F, S_F)^l$$

where k and l are non-negative integers such that $k + l$ is greater than 0 and (α_G, τ_G, S_G) and (α_F, τ_F, S_F) are the prime homotopy data triples defined in Example 4.3.

The following lemma extends Turaev's Lemma 3.3.1 in [15] to the case of general S .

Lemma 4.9. *Let $(\alpha, \tau_\alpha, S_\alpha)$ be a homotopy data triple and let $(\beta, \tau_\beta, S_\beta)$ be a factor of $(\alpha, \tau_\alpha, S_\alpha)$. We denote the equivalence relation given by $(\alpha, \tau_\alpha, S_\alpha)$ as $\sim_\alpha$*

and the equivalence relation given by $(\beta, \tau_\beta, S_\beta)$ as $\sim_\beta$. Now let p and p' be two nanophrases in $\mathcal{P}(\beta)$. If $p \sim_\alpha p'$, then $p \sim_\beta p'$.

Proof. This is proved in the same way as Lemma 3.3.1 of [15]. $\square$

Remark 4.10. In Lemma 3.3.1 of [15], Turaev proved the case where S_α is diagonal and p and p' are étale words. An étale word is a word on an α -alphabet, so all nanowords are étale words (see [15] for more details).

5. AN INVARIANT FOR NANOWORDS

In this section we fix a composite homotopy data triple (α, τ, S) and let its prime factors be denoted by (α_i, τ_i, S_i) for i running from 1 to k for some k greater than 1. Let $\sim$ be the homotopy given by (α, τ, S) and, for each i , let $\sim_i$ be the homotopy given by (α_i, τ_i, S_i) .

For any positive integer n , we say that a map from $\hat{n}$ to $\hat{k}$ is *locally variable* if there does not exist an i for which $\theta(i)$ equals $\theta(i+1)$. Let $\mathcal{P}_A(\alpha)$ be the set of pairs (p, θ) , where p is a nanophrase over α and θ is a locally variable map from $\widehat{\text{nc}(p)}$ to $\hat{k}$. We extend the definition of isomorphism and homotopy moves of nanophrases to $\mathcal{P}_A(\alpha)$ by saying that $(p, \theta) \sim (p', \theta')$ if $p \sim p'$.

Let $\mathcal{P}_R(\alpha)$ be the subset of $\mathcal{P}_A(\alpha)$ consisting of pairs (p, θ) for which every letter X appearing in the i th component of p , $|X|$ is in $\alpha_{\theta(i)}$, for all i . Note that if p is a nanophrase over α with no empty components then if (p, θ) and (p, θ') are both in $\mathcal{P}_R(\alpha)$ we can conclude that θ and θ' are equal.

Note that we can define a map Γ from $\mathcal{P}_A(\alpha)$ to $\mathcal{P}_R(\alpha)$ as follows. Given (p, θ) in $\mathcal{P}_A(\alpha)$ we derive p' from p by deleting all letters X which have projections not matching θ . In other words, for each i , we delete every letter X in the i th component of p for which $|X|$ is not in $\alpha_{\theta(i)}$.

We define an equivalence relation on $\mathcal{P}_R(\alpha)$, written $\sim_K$, as follows. Let (p, θ) and (p', θ) be elements of $\mathcal{P}_R(\alpha)$. Then $(p, \theta) \sim_K (p', \theta)$ if there exists a sequence of elements (p_i, θ) of $\mathcal{P}_R(\alpha)$, for i running from 0 to r for some r , such that p_0 is p , p_r is p' and p_i is related to p_{i+1} by a single homotopy move or an isomorphism.

Note that $(p, \theta) \sim_K (p', \theta)$ implies $(p, \theta) \sim (p', \theta)$. The following lemmas show that $\sim_K$ can be viewed as a restriction of $\sim$ to $\mathcal{P}_R(\alpha)$.

Lemma 5.1. *Let (p, θ) and (p', θ) be two elements in $\mathcal{P}_A(\alpha)$. If $(p, \theta) \sim (p', \theta)$ then $\Gamma((p, \theta)) \sim_K \Gamma((p', \theta))$.*

Proof. This can be proved in a similar way to Lemma 3.2. $\square$

Lemma 5.2. *Let (p, θ) and (p', θ) be two elements in $\mathcal{P}_R(\alpha)$. If $(p, \theta) \sim (p', \theta)$ then $(p, \theta) \sim_K (p', \theta)$.*

Proof. This can be proved in a similar way to Lemma 3.3. $\square$

Let (p, θ) be an element of $\mathcal{P}_R(\alpha)$. We fix an integer i and let O_i be the subset of $\widehat{\text{nc}(p)}$ consisting of elements j where $\theta(j)$ is equal to i . Then we define $s_i((p, \theta))$ to be $x(p, O_i)$ (which was defined in Section 3). Note that $s_i((p, \theta))$ is the subnanophrase of p consisting of the components containing letters which map to α_i and so $s_i((p, \theta))$ is a nanophrase over α_i .

Proposition 5.3. *Let (p, θ) and (p', θ) be two elements of $\mathcal{P}_R(\alpha)$. Then $(p, \theta) \sim_K (p', \theta)$ if and only if $s_i((p, \theta)) \sim_i s_i((p', \theta))$ for all i .*

Proof. We first assume that $s_i((p, \theta)) \sim_i s_i((p', \theta))$ for all i . Note that each component of p appears as a component in exactly one of the nanophrases $s_i((p, \theta))$. Since $s_1((p, \theta)) \sim_1 s_1((p', \theta))$, there is a sequence of homotopy moves going

from $s_1((p, \theta))$ to $s_1((p', \theta))$ which can be applied directly to the corresponding components in (p, θ) . Call the result (p_1, θ) . Then $s_1(p_1, \theta)$ is isomorphic to $s_1((p', \theta))$ and $s_j(p_1, \theta)$ is equal to $s_j(p, \theta)$ for all j not equal to 1. We also have $(p_1, \theta) \sim_K (p, \theta)$. Starting with (p_1, θ) we repeat the process, this time using the fact that $s_2((p, \theta)) \sim_2 s_2((p', \theta))$. The result is (p_2, θ) which is $\sim_K$ equivalent to (p, θ) . By repeating the process for each i , the final result (p_k, θ) is $\sim_K$ equivalent to (p, θ) and isomorphic to (p', θ) . Thus we have shown that $(p, \theta) \sim_K (p', \theta)$.

We now assume that $(p, \theta) \sim_K (p', \theta)$. This implies that $p \sim p'$. Now, fixing an integer i , let O_i be the subset of $\text{nc}(p)$ consisting of elements j where $\theta(j)$ is equal to i . Then, by definition, $s_i((p, \theta))$ is $x(p, O_i)$ and $s_i((p', \theta))$ is $x(p', O_i)$. Since $p \sim p'$, Lemma 3.7 implies $x(p, O_i) \sim x(p', O_i)$ and thus, using Lemma 4.9, $x(p, O_i) \sim_i x(p', O_i)$. $\square$

We define some new moves on elements of $\mathcal{P}_R(\alpha)$ which allow us to remove or add empty components. Let (p, θ) be an element of $\mathcal{P}_R(\alpha)$. We write n for $\text{nc}(p)$ and w_j for the j th component of p . If w_i is $\emptyset$ we can apply a simple reduction or concatenating reduction which are defined as follows.

Simple Reduction: if $i = 1$, $i = n$ or $\theta(i - 1) \neq \theta(i + 1)$,

$$(w_1 | \dots | w_{i-1} | \emptyset | w_{i+1} | \dots | w_n, \theta) \longrightarrow (w_1 | \dots | w_{i-1} | w_{i+1} | \dots | w_n, \theta')$$

where θ' is a map from $\widehat{n-1}$ to $\hat{k}$ defined by

$$\theta'(j) = \begin{cases} \theta(j) & \text{if } j < i \\ \theta(j+1) & \text{if } j \geq i. \end{cases}$$

Note that if n is 1 and w_1 is $\emptyset$, applying a simple reduction gives the pair $(\emptyset_P, \theta_\emptyset)$ where $\theta_\emptyset$ is the empty map (from $\emptyset$ to $\hat{k}$). The inverse of a simple reduction is called a *simple augmentation*. It allows us to insert a new empty component into p either between two components or at one of the ends, as long as we can make a corresponding change to θ to get a locally variable map.

Concatenating Reduction: if $i \neq 1$, $i \neq n$ and $\theta(i - 1) = \theta(i + 1)$,

$$(w_1 | \dots | w_{i-1} | \emptyset | w_{i+1} | \dots | w_n, \theta) \longrightarrow (w_1 | \dots | w_{i-1} w_{i+1} | \dots | w_n, \theta')$$

where θ' is a map from $\widehat{n-2}$ to $\hat{k}$ defined by

$$\theta'(j) = \begin{cases} \theta(j) & \text{if } j < i \\ \theta(j+2) & \text{if } j \geq i. \end{cases}$$

The inverse of this move is called a *splitting augmentation*. It allows us to split a component of p into two parts (where possibly one or both may be empty), as long as we can make a corresponding change to θ to get a locally variable map.

Collectively we refer to simple reductions and concatenating reductions as *reductions* and simple augmentations and splitting augmentations as *augmentations*.

Remark 5.4. Suppose (p, θ) is an element of $\mathcal{P}_R(\alpha)$ such that the i th component of p is empty. Then by the conditions on i and θ given in the definitions of the reduction moves, there is one and only one way that we can apply a reduction to (p, θ) to remove the i th component. Thus we can unambiguously refer to this reduction as *the reduction of the i th component* of (p, θ) .

Let $\sim_P$ be the equivalence relation on $\mathcal{P}_R(\alpha)$ generated by $\sim_K$, reductions and augmentations. We can define a map Ω from $\mathcal{P}_R(\alpha)$ to $\mathcal{N}(\alpha)$ by mapping (p, θ) to $\chi(p)$ (recall that χ is the concatenating map defined in Section 3). Then we have the following lemma.

Lemma 5.5. *Let (p, θ) and (p', θ') be elements of $\mathcal{P}_R(\alpha)$. If $(p, \theta) \sim_P (p', \theta')$ then $\Omega((p, \theta)) \sim \Omega((p', \theta'))$.*

Proof. If $(p, \theta) \sim_K (p', \theta')$, then θ and θ' are equal and $p \sim p'$. Then by Lemma 3.1, $\chi(p) \sim \chi(p')$ and so $\Omega((p, \theta)) \sim \Omega((p', \theta'))$. So, to prove the result, it is enough to prove the case where (p', θ') is derived from (p, θ) by a single reduction. However in this case, it is easy to see that $\chi(p)$ and $\chi(p')$ must be equal and so $\Omega((p, \theta)) \sim \Omega((p', \theta'))$. $\square$

We define a map ψ from $\mathcal{N}(\alpha)$ to $\mathcal{P}_R(\alpha)$ as follows. Let w be a nanoword in $\mathcal{N}(\alpha)$. If w is $\emptyset$ then we map w to $\emptyset_P$. Otherwise, there is a unique integer $r > 0$, a unique sequence of words $w_1, w_2, \dots, w_r$ and a unique map θ from $\hat{r}$ to $\hat{k}$ such that

- (1) each w_i is not $\emptyset$;
- (2) w is equal to $w_1 w_2 \dots w_r$;
- (3) for all i , if X appears in w_i , $|X|$ is in $\alpha_{\theta(i)}$;
- (4) for any i from 1 to $r - 1$, $\theta(i)$ is not equal to $\theta(i + 1)$.

Define p to be the nanophrase $w_1 |w_2| \dots |w_r$ where the letters in p have the same projection as they do in w . Then (p, θ) is in $\mathcal{P}_R(\alpha)$. We define $\psi(w)$ to be (p, θ) .

Example 5.6. Let α be the set $\{a, b\}$. Suppose that under factorization it factors into two sets, $\{a\}$, which we call α_1 , and $\{b\}$, which we call α_2 . Then k is 2.

Let $\mathcal{A}$ be the α -alphabet $\{A, B, C, D, E\}$ with projection given by $|A| = |B| = |C| = a$ and $|D| = |E| = b$. Now consider the nanoword $(\mathcal{A}, w)$ where w is $ABCBDCAEDE$.

Then w_1 is $ABC B$, w_2 is D , w_3 is $C A$ and w_4 is $E D E$. So n is 4 and θ is a map from $\hat{4}$ to $\hat{2}$ which maps 1 and 3 to 1 and maps 2 and 4 to 2. Thus ψ maps $(\mathcal{A}, w)$ to $((\mathcal{A}, w_1 |w_2| w_3 |w_4), \theta)$.

Note that the image of ψ consists of all elements (p, θ) in $\mathcal{P}_R(\alpha)$ such that p has no empty components.

Lemma 5.7. *Let w and w' be elements of $\mathcal{N}(\alpha)$. If $w \sim w'$, then $\psi(w) \sim_P \psi(w')$.*

Proof. It is enough to prove the case where w and w' are related by an isomorphism or a single homotopy move.

If w and w' are related by an isomorphism, then the nanophrases in $\psi(w)$ and $\psi(w')$ are isomorphic and so $\psi(w) \sim_P \psi(w')$.

We now assume w and w' are related by a single homotopy move. We write $\psi(w)$ as (p, θ) and $\psi(w')$ as (p', θ') .

Suppose w and w' are related by an H1 move. If θ and θ' are equal, then p is related to p' by a corresponding H1 move. If not, we assume that w' is derived from w by removing a letter. By making the corresponding H1 move on (p, θ) and then applying an appropriate reduction we can derive (p', θ') . Thus, in either case, $\psi(w) \sim_P \psi(w')$.

Suppose w and w' are related by an H2 move. If θ and θ' are equal, then p is related to p' by a corresponding H2 move. If not, we assume that w' is derived from w by removing two letters. By making the corresponding H2 move on (p, θ) and then applying appropriate reductions we can derive (p', θ') . Thus, in either case, $\psi(w) \sim_P \psi(w')$.

If w and w' are related by an H3 move, then θ and θ' are equal and p is related to p' by a corresponding H3 move. Thus $\psi(w) \sim_P \psi(w')$. $\square$

Theorem 5.8. *There is a bijection between the homotopy classes of $\mathcal{N}(\alpha)$ and the equivalence classes of $\mathcal{P}_R(\alpha)$ under $\sim_P$. In other words*

$$\mathcal{N}(\alpha) / \sim \cong \mathcal{P}_R(\alpha) / \sim_P .$$

Proof. The map ψ induces a map ψ_P from $\mathcal{N}(\alpha)/\sim$ to $\mathcal{P}_R(\alpha)/\sim_P$, which by Lemma 5.7 is well-defined. Note that for any nanoword w in $\mathcal{N}(\alpha)$, $\Omega(\psi(w))$ is w . Then by Lemma 5.5, ψ_P is bijective. $\square$

Let $\mathcal{K}(\alpha)$ be the set of equivalence classes of $\mathcal{P}_R(\alpha)$ under $\sim_K$. Let c be an element of $\mathcal{K}(\alpha)$ and (p, θ) be an element in c . We define $\text{nc}(c)$ to be $\text{nc}(p)$. Note that if (q, θ) is another element in c then $\text{nc}(p)$ is equal to $\text{nc}(q)$, and so $\text{nc}(c)$ is well-defined.

We say that an element c of $\mathcal{K}(\alpha)$ is *i-reducible* if there exists a pair (p, θ) in c such that the i th component of p is $\emptyset$.

Lemma 5.9. *Let c be an element of $\mathcal{K}(\alpha)$. Suppose (p, θ) and (p', θ') are elements in c such that p and p' are both *i-reducible*. Then there exists a sequence of elements of c going from (p, θ) to (p', θ') such that the nanophrases in each consecutive pair are related by a single homotopy move or isomorphism and each nanophrase in the sequence is *i-reducible*.*

Proof. This is proved in a similar way to Lemma 3.3. $\square$

We note that the equivalence relation $\sim_P$ on the set $\mathcal{P}_R(\alpha)$ induces an equivalence relation on $\mathcal{K}(\alpha)$ which we also write $\sim_P$.

Let (p', θ') be the result the reduction of the i th component of (p, θ) and let c' be the element of $\mathcal{K}(\alpha)$ which contains (p', θ') . We say that c' is the *i-reduction* of c . The following lemma shows that this concept is well-defined.

Lemma 5.10. *Let c be an element of $\mathcal{K}(\alpha)$. If c' and c'' are *i-reductions* of c , c' equals c'' .*

Proof. As c' is an *i-reduction* of c , there exists (p, θ) in c such that the i th component of p is $\emptyset$ and the reduction of the i th component of (p, θ) gives (p', θ') in c' . Similarly, as c'' is an *i-reduction* of c , there exists (q, θ) in c such that the i th component of q is $\emptyset$ and the reduction of the i th component of (q, θ) gives (q', θ'') in c'' . By Lemma 5.9 there exists a sequence of nanophrases p_j going from p to q such that each (p_j, θ) is in c and the i th component of p_j is $\emptyset$. Then for each j it is easy to check that the reduction of the i th component of (p_j, θ) is equivalent, under $\sim_K$ to the reduction of the i th component of (p_{j+1}, θ) . Thus $(p', \theta') \sim_K (q', \theta'')$ and so c' equals c'' . $\square$

We say that an element c of $\mathcal{K}(\alpha)$ is *reducible* if it is *i-reducible* for some i . An element c of $\mathcal{K}(\alpha)$ is *irreducible* if it is not reducible.

Lemma 5.11. *Let c be an element of $\mathcal{K}(\alpha)$ and let i and j be different integers in $\widehat{\text{nc}(c)}$. If c is both *i-reducible* and *j-reducible*, then there exists an element (p, θ) in c such that both the i th and j th components of p are $\emptyset$.*

Proof. As c is *i-reducible*, there exists an element (q, θ) in c such that the i th component of q is $\emptyset$. As c is *j-reducible*, there exists an element (q', θ) in c such that the j th component of q' is $\emptyset$. Recall that for a nanophrase p and a subset O of $\widehat{\text{nc}(p)}$, f_O maps p to the nanophrase derived from p by deleting all letters that appear at least once in any component with index appearing in O . Let O be the set $\{j\}$. As $(q, \theta) \sim_K (q', \theta)$, by a similar argument to the proof of Lemma 3.4, we can show that $(q, \theta) \sim_K (f_O(q), \theta)$. Thus $(f_O(q), \theta)$ is in c and both the i th and j th components of $f_O(q)$ are $\emptyset$. $\square$

Lemma 5.12. *Let c be an element of $\mathcal{K}(\alpha)$ and let i and j be integers in $\widehat{\text{nc}(c)}$ with i less than j . Suppose c is both *i-reducible* and *j-reducible*. Let c_i be the result of the *i-reduction* of c and let c_j be the result of the *j-reduction* of c . Then one of the following cases holds.*

- (1) c_i is equal to c_j ;
- (2) c_i is a reduction of c_j ;
- (3) c_j is a reduction of c_i ;
- (4) there exists an element c' of $\mathcal{K}(\alpha)$ which is a reduction both of c_i and c_j .

Proof. By Lemma 5.11 we know that c contains an element (p, θ) such that the i th and j th components of p are $\emptyset$. We write (p_i, θ_i) for the i -reduction of p and (p_j, θ_j) for the j -reduction of p . Then (p_i, θ_i) is in c_i and (p_j, θ_j) is in c_j .

Assume that j is greater than $i + 2$. Then the set of components involved in the i -reduction and the set of components involved in the j -reduction have no components in common. So (p_j, θ_j) will be i -reducible and the i -reduction will be of the same type as the i -reduction of (p, θ) . Similarly, there will be a reduction available in (p_i, θ_i) corresponding to the j -reduction of (p, θ) . If the i -reduction of (p, θ) is simple, (p_i, θ_i) will be $(j - 1)$ -reducible. If the i -reduction of (p, θ) is concatenating, (p_i, θ_i) will be $(j - 2)$ -reducible. It is easy to check that in all cases, applying the two reductions in either order gives the same result.

Let (p', θ') be the result of applying an i -reduction to (p_j, θ_j) and let c' be the equivalence class of (p', θ') under $\sim_K$. Then as c' is a reduction of both c_i and c_j , Case 4 in the statement of the lemma holds.

For example, the case where both reductions are simple is shown in Figure 1. In the figure w_t is the t th component of p for each t and labels on the arrows indicate which component is being reduced.




FIGURE 1. Simple reductions





Now assume that j is $i + 2$. If the i -reduction or j -reduction of (p, θ) is simple then the situation is similar to the case where j is greater than $i + 2$ and so Case 4 in the statement of the lemma holds. The case where both the i -reduction and the j -reduction of (p, θ) are concatenating is shown in Figure 2. Note that because both reductions are concatenating, $\theta(i - 1)$, $\theta(i + 1)$ and $\theta(i + 3)$ are all equal. Thus, again, Case 4 in the statement of the lemma holds.

Now assume that j is $i + 1$. We consider cases based on the types of the reductions of (p, θ) .

If both the i -reduction and the j -reduction are concatenating then the results of the reductions are equivalent, as shown in Figure 3. Thus Case 1 in the statement of the lemma holds.

The case where the i -reduction is simple and the j -reduction is concatenating is shown in Figure 4. Thus Case 2 in the statement of the lemma holds.

The case where the i -reduction is concatenating and the j -reduction is simple is shown in Figure 5. Thus Case 3 in the statement of the lemma holds.

FIGURE 2. Concatenating reductions when $j = i + 2$ FIGURE 3. Concatenating reductions when $j = i + 1$ FIGURE 4. Simple i -reduction, concatenating j -reduction when $j = i + 1$ FIGURE 5. Concatenating i -reduction, simple j -reduction when $j = i + 1$

The case where both the i -reduction and the j -reduction are simple splits into two subcases. The first subcase is when i is 1 or j is n or $\theta(i-1)$ is not equal to $\theta(i+2)$. Then (p_i, θ_i) is i -reducible by a simple reduction and (p_j, θ_j) is also

i -reducible by a simple reduction. Either reduction gives the same result and so Case 4 in the statement of the lemma holds.

The second subcase is when i is not 1, j is not n and $\theta(i-1)$ is equal to $\theta(i+2)$. Then both (p_i, θ_i) and (p_j, θ_j) are i -reducible by concatenating reductions as shown in Figure 6. Note that in this case p_i is equal to p_j , but as $\theta(i)$ is not equal to $\theta(i+1)$, θ_i and θ_j are different. From the figure we see that Case 4 in the statement of the lemma holds. $\square$




FIGURE 6. Simple reductions when $j = i + 1$ and $\theta(i-1) = \theta(i+2)$

Let F be a set with an equivalence relation defined on it in terms of two types of moves, ‘reducing’ moves and the inverses of these moves. An element of F is said to be *reduced* if it does not admit a reducing move. A *reducing chain* of elements of F is a sequence $g_1, g_2, \dots, g_n$ such that g_{i+1} is the result of a reducing move applied to g_i for all i .

The following lemma, known as the Diamond Lemma was proved by Newman in [12]. The version we give here is based on the version of the lemma appearing in [1, page 26].

Lemma 5.13 (Diamond Lemma, Newman). *Let F be a set as above, such that the following conditions are satisfied:*

- (1) *Finiteness condition. For each element g in F there exists an integer r (depending on g) such that no reducing chain starting from g has more than r terms.*
- (2) *Confluence condition. If an element g in F can be transformed to g_1 by one reducing move and g_2 by another, then there exists an element g' such that, for each i , either g' is equal to g_i or g_i can be transformed into g' by applying one or more reducing moves.*

Then each equivalence class of F contains exactly one reduced element.

Proposition 5.14. *Each equivalence class of $\mathcal{K}(\alpha)$ under $\sim_P$ contains exactly one reduced element.*

Proof. We prove this using Lemma 5.13. We just need to show that the Confluence condition and the Finiteness condition hold for $\mathcal{K}(\alpha)$.

The fact that the Confluence condition holds is shown by Lemma 5.12. On the other hand, a reducing chain starting from c , an element of $\mathcal{K}(\alpha)$, can have no more than $\text{nc}(c) + 1$ elements, because a reduction move reduces the number of components by at least 1. Thus the Finiteness condition of Lemma 5.13 also holds. $\square$

We define a map R from $\mathcal{K}(\alpha)$ to itself as follows. For an element c in $\mathcal{K}(\alpha)$, we define $R(c)$ to be the reduced element in the equivalence class of c under $\sim_P$. By Proposition 5.14, the map R is well-defined.

Assuming that we can tell whether any element of $\mathcal{K}(\alpha)$ is reducible or not (which in general is a difficult problem), it is easy to calculate $R(c)$ for any c using an inductive process. If c is irreducible, $R(c)$ is c . If not, c can be reduced to some other element of $\mathcal{K}(\alpha)$, say c' and $R(c)$ is equal to $R(c')$. As was shown in the proof of Proposition 5.14, the Finiteness condition holds, so this process will terminate in a finite number of steps.

Let w be a nanoword in $\mathcal{N}(\alpha)$. Then $\psi(w)$ is an element in $\mathcal{P}_R(\alpha)$. Let c be the element of $\mathcal{K}(\alpha)$ which contains $\psi(w)$. Then we define $R_K(w)$ to be $R(c)$.

Theorem 5.15. *For a nanoword w in $\mathcal{N}(\alpha)$, $R_K(w)$ is a homotopy invariant of w .*

Proof. This follows from Theorem 5.8 and Proposition 5.14. $\square$

In particular, this means that $\text{nc}(R_K(w))$ and the θ associated with $R_K(w)$ are invariants of w . We write $c_R(w)$ for $\text{nc}(R_K(w))$ and write $\theta_R(w)$ for the θ associated with $R_K(w)$. We also define $P_{R,i}(w)$ to be $s_i((p, \theta))$ where (p, θ) is an element in $R_K(w)$. The nanophrase $s_i((p, \theta))$ is a nanophrase in $\mathcal{P}(\alpha_i)$ and is, modulo $\sim_i$, an invariant of w .

We have the following corollary of Theorem 5.15 which show that $\theta_R(w)$ and the sequence of nanophrases $P_{R,i}(w)$ together give a complete invariant for w .

Corollary 5.16. *Let w and w' be nanowords over α . Then $w \sim w'$ if and only if $\theta_R(w)$ and $\theta_R(w')$ are equal and, for all i , $P_{R,i}(w) \sim_i P_{R,i}(w')$*

Proof. This follows from Theorem 5.15 and Proposition 5.3. $\square$

We say that w is *c-minimal* if $\text{nc}(\psi(w))$ is equal to $c_R(w)$. The following proposition shows how the property of being minimal and the property of being c-minimal are related.

Proposition 5.17. *Let w be a nanoword. If w is minimal, then it is c-minimal.*

Proof. We write (p, θ) for $\psi(w)$ and assume that w is minimal but not c-minimal for a contradiction. The assumption that w is not c-minimal is equivalent to assuming that $\psi(w)$ is not in $R_K(w)$. This means that $\psi(w)$ is i -reducible for some i . Thus $(p, \theta) \sim_K (p', \theta)$ for some nanophrase p' for which the i th component is empty.

Let O equal $\{i\}$. Then by Lemma 3.4, $p \sim f_O(p)$ and so $(p, \theta) \sim_K (f_O(p), \theta)$. Note that because (p, θ) is in the image of ψ , the i th component of p is not empty. Thus $\text{rank}(p)$ is greater than $\text{rank}(f_O(p))$.

Let (q, θ') be the result of applying an i -reduction to $(f_O(p), \theta)$. Then $\text{rank}(q)$ is equal to $\text{rank}(f_O(p))$ and so less than $\text{rank}(p)$. As $(q, \theta') \sim_P (p, \theta)$, by Lemma 5.5, $\Omega((q, \theta')) \sim \Omega((p, \theta))$. Now $\Omega((p, \theta))$ is w , so $\Omega((q, \theta')) \sim w$. However, $\text{rank}(\Omega((q, \theta')))$ is equal to $\text{rank}(q)$ which is less than $\text{rank}(p)$, so $\text{hr}(w)$ is less than $\text{rank}(w)$ contradicting the fact that w is minimal. $\square$

Remark 5.18. Note that if a nanoword w is c-minimal it does not necessarily mean that it is minimal. For example, suppose w is a non-trivial minimal nanoword. Then w has the form $XYXz$ for some letter X and some words y and z . Then the nanoword $AAXyXz$ where $|A|$ is equal to $|X|$ is clearly c-minimal but not minimal.

Theorem 5.19. *Let (α, τ, S) be a composite homotopy data triple and let its prime factors be denoted by (α_i, τ_i, S_i) . Let w be a nanoword over α . Then*

$$(5.1) \quad \text{hr}(w) = \sum_i \text{hr}(P_{R,i}(w)).$$

Proof. Let u be a minimal nanoword homotopic to w . Then $\text{rank}(u)$ is equal to $\text{hr}(w)$. By Proposition 5.17 $\psi(u)$ is in $R_K(w)$. Then $\text{hr}(P_{R,i}(u))$ is equal to $\text{hr}(P_{R,i}(w))$ and clearly

$$(5.2) \quad \text{hr}(u) \geq \sum_i \text{hr}(P_{R,i}(w)).$$

On the other hand, for each i , let q_i be a minimal nanophrase such that $q_i \sim_i P_{R,i}(w)$. Then let $(q, \theta_R(w))$ be the element of $\mathcal{P}_R(\alpha)$ such that $s_i((q, \theta_R(w)))$ is equal to q_i for all i . Then $(q, \theta_R(w))$ is in $R_K(w)$. Then by Lemma 5.5, $\Omega((q, \theta_R(w))) \sim u$ and so

$$(5.3) \quad \text{hr}(u) \leq \sum_i \text{hr}(P_{R,i}(w)).$$

Combining the inequalities (5.2) and (5.3) gives (5.1). $\square$

Remark 5.20. If (α, τ, S) is a prime homotopy data triple then the results of this section hold, but are trivial. The set $\mathcal{P}_R(\alpha)$ consists of the element $(\emptyset_P, \theta_\emptyset)$, where $\theta_\emptyset$ is the empty map, and elements $(p, \theta : 1 \mapsto 1)$ where p is a 1-component nanophrase. Thus $\theta_R(w)$ can take one of two values depending on whether w is homotopic to $\emptyset$ or not. However, to calculate the value of $\theta_R(w)$ for a given nanoword w we must show whether w is homotopic to $\emptyset$ or not and so the invariant is trivial. Similarly, if w is not homotopic to $\emptyset$, $P_{R,1}(\psi(w))$ is given by w , and again the invariant is trivial. In this case, Theorem 5.19 is reduced to the tautological statement that $\text{hr}(w)$ is equal to itself.

We finish this section by briefly considering some symmetries of nanowords defined by Turaev in [15]. We start by recalling some definitions. Given a nanoword w , the *opposite nanoword* of w is the nanoword given by taking the letters of w in the reverse order and preserving the projection of the letters. We write w^- for the opposite nanoword of w . The *inverse nanoword* of w , written $\overline{w}$, has the same Gauss word as w but the projections of the letters are composed with τ . In other words, for all A appearing in w , $|A|_{\overline{w}}$, the projection of A in $\overline{w}$, is equal to $\tau(|A|_w)$, where $|A|_w$ is the projection of A in w .

We extend these definitions to nanophrases. Given a nanophrase p , the *opposite nanophrase* of p , written p^- is the nanophrase given by taking the components in reverse order and, for each component, reversing the order of the letters in the component. The *inverse nanophrase* of p , written $\overline{p}$, has the same Gauss phrase as p but the projections of the letters are composed with τ . These operations are involutions, so the opposite nanophrase of p^- is p and the inverse nanophrase of $\overline{p}$ is p . The two operations commute. Note that for a 1-component nanophrase, which we can consider to be a nanoword, the definitions given for nanowords and nanophrases coincide.

Example 5.21. Let α be the set $\{a, b\}$ and let τ be the map swapping a with b . Let p be the nanophrase $ABC|AC|B$ where $|A|$ and $|C|$ are a and $|B|$ is b . Then p^- is $B|CA|CBA$ where the letters have the same projections as p . The nanophrase $\overline{p}$ is $ABC|AC|B$ where $|A|$ and $|C|$ are b and $|B|$ is a .

In [15], Turaev defined the following terms. Given a homotopy, a nanoword w is *homotopically symmetric* (with respect to that homotopy) if w is homotopic to w^- . Given a homotopy, a nanoword w is *homotopically skew-symmetric* (with respect to that homotopy) if w is homotopic to $(\overline{w})^-$.

We fix a homotopy data triple (α, τ, S) . Let κ_τ be the involution $\tau \times \tau \times \tau$ on $\alpha \times \alpha \times \alpha$, and let κ_i be the involution which maps (a, b, c) to (c, b, a) on $\alpha \times \alpha \times \alpha$.

Suppose S is invariant under κ_τ . Then Turaev observed in [15] that if u and v are homotopic under the homotopy given by (α, τ, S) , then $\overline{u}$ and $\overline{v}$ are also homotopic under the same homotopy. Then it is easy to see that, for any nanoword w , $c_R(\overline{w})$ is equal to $c_R(w)$, $\theta_R(\overline{w})$ is equal to $\theta_R(w)$ and, for all i ,

$$P_{R,i}(\overline{w}) = \overline{P_{R,i}(w)}.$$

Now suppose that S is invariant under κ_i . Then Turaev observed in [15] that if u and v are homotopic under the homotopy given by (α, τ, S) , then u^- and v^- are also homotopic under the same homotopy. Again, it is easy to see that, for any nanoword w , $c_R(w^-)$ is equal to $c_R(w)$, $\theta_R(w^-)(i)$ is equal to $\theta_R(w)(c_R(w) + 1 - i)$ for all i and

$$P_{R,i}(w^-) = (P_{R,i}(w))^-.$$

for all i .

So if S is invariant under κ_i , w can only be homotopically symmetric if

$$(5.4) \quad (\theta_R(w))(i) = \theta_R(w)(c_R(w) + 1 - i)$$

for all i . As $\theta_R(w)$ is locally variable, this implies that $c_R(w)$ must be odd. Thus if $c_R(w)$ is even or (5.4) does not hold for all i , w cannot be homotopically symmetric. Similarly, if S is invariant under κ_i and κ_τ , w cannot be homotopically skew-symmetric if $c_R(w)$ is even or (5.4) does not hold for all i .

6. DECIDABILITY

We say that a homotopy is *reduction decidable* if we have a finite time algorithm which, for any nanophrase p and any integer i , determines whether p is i -reducible or not. We say that a homotopy is *equality decidable* if we have a finite time algorithm which, for any two nanophrases p and q , determines whether p is equivalent to q or not.

Proposition 6.1. *Let (α, τ, S) be a composite homotopy data triple and let its prime factors be denoted by (α_i, τ_i, S_i) . Suppose, for each i , the homotopy given by (α_i, τ_i, S_i) is reduction decidable. Then there is a finite time algorithm which, for any nanoword over α , w , produces a nanoword $R(w)$ such that $\psi(R(w))$ is in $R_K(w)$.*

Proof. We give a finite time algorithm to produce $R(w)$ from an arbitrary nanoword w .

We first note that $\psi(w)$ is reducible if and only if $\psi_i(w)$ is reducible for some i . As, for each i , the homotopy given by (α_i, τ_i, S_i) is reduction decidable, we can determine whether or not $\psi(w)$ is reducible in finite time.

If $\psi(w)$ is not reducible then $\psi(w)$ is in $R_K(w)$ and we define $R(w)$ to be w .

If $\psi(w)$ is reducible, then it is j -reducible for some j . Let O be the subset of $\widehat{\text{nc}(p)}$ consisting of the single element j . Writing (p, θ) for $\psi(w)$, let p' be $f_O(\psi(w))$. Then by Lemma 3.4 and Lemma 5.2, $(p, \theta) \sim_K (p', \theta)$. Let c be the equivalence class under $\sim_K$ which contains $\psi(w)$ and let c' be the equivalence class under $\sim_K$ which contains $\psi(\chi(p'))$. It is clear that c' is derivable from c by a series of one or more reductions. We define $R(w)$ to be $R(\chi(p'))$. This definition is recursive, but the recursion will terminate because $\text{nc}(\chi(p'))$ is less than $\text{nc}(w)$. $\square$

We note that $R(w)$ is not necessarily well-defined as it depends on the order that the reductions are made. For example, consider the nanoword $ACADDBBC$ where $|A|$ and $|B|$ are in α_1 and $|C|$ and $|D|$ are in α_2 . Then $\psi(w)$ is $(A|C|A|DD|BB|C, \theta)$ where θ maps odd numbers to 1 and even numbers to 2. The 4th and 5th components of $\psi(w)$ are clearly reducible by an H1 move. If we apply the above process to the 4th component we get $(A|C|ABB|C, \theta')$ where θ' maps odd numbers to 1

and even numbers to 2. If we apply the above process to the 5th component we get $(A|C|A|DDC, \theta')$. Using the linking matrix described in Section 7.2 we can show that both these elements are reduced. Thus $R(w)$ can be $(A|C|ABB|C, \theta')$ or $(A|C|A|DDC, \theta')$ depending on the order we make the reductions.

Theorem 6.2. *Let (α, τ, S) be a composite homotopy data triple and let its prime factors be denoted by (α_i, τ_i, S_i) . Suppose, for each i , the homotopy given by (α_i, τ_i, S_i) is reduction decidable and equality decidable. Then, there is a finite time algorithm which for any two nanowords over α , w and w' , determines whether or not they are equivalent under the homotopy given by (α, τ, S) .*

Proof. Let w and w' be two nanowords in α . By Proposition 6.1 we can calculate $R(w)$ and $R(w')$ in finite time. We write (p, θ) for $R(w)$ and (p', θ') for $R(w')$. Now, as $\psi(R(w))$ and $\psi(R(w'))$ are irreducible, if θ and θ' are not equal, w and w' are not equivalent.

If θ and θ' are equal, then by Corollary 5.16, $\psi(w) \sim \psi(w')$ if and only if $P_{R,i}(w) \sim_i P_{R,i}(w')$ for all i . Now, for all i , $P_{R,i}(w) \sim_i P_{R,i}(w')$ can be determined in finite time because $\sim_i$ is equality decidable. $\square$

Remark 6.3. Recall the definitions of (α_G, τ_G, S_G) and (α_F, τ_F, S_F) from Example 4.3. By Theorem 6.2 and Example 4.8 we may conclude the following. If (α_G, τ_G, S_G) and (α_F, τ_F, S_F) are both reduction and equality decidable then for any S diagonal homotopy there is a finite time algorithm which can determine homotopy equivalence of arbitrary pairs of nanowords. Whether (α_G, τ_G, S_G) or (α_F, τ_F, S_F) are reduction or equality decidable is an open question.

We say that a homotopy has a *normal form* if there exists a finite time algorithm taking a nanophrase p and producing a nanophrase $n(p)$ which satisfies the following conditions:

- (1) $n(p) \sim p$ for all p ,
- (2) if $p \sim p'$, then $n(p)$ equals $n(p')$ for all p and p' .

If, for all p , $\text{rank}(n(p))$ is equal to $\text{hr}(p)$, then we say that n is a *minimizing normal form*.

Proposition 6.4. *If a homotopy has a minimizing normal form n , it is reduction decidable and equality decidable.*

Proof. The fact that the homotopy is equality decidable immediately follows from the existence of a normal form (minimizing or not). Now p is i -reducible if and only if $n(p)$ is i -reducible. If the i th component of $n(p)$ is $\emptyset$, then $n(p)$ is i -reducible by definition. On the other hand, suppose that the i th component of $n(p)$ is not $\emptyset$. Then as n is a minimizing normal form, $\text{rank}(n(p))$ is equal to $\text{hr}(p)$ and so by Lemma 3.5, $n(p)$ is not i -reducible. Thus p is i -reducible if and only if the i th component of $n(p)$ is trivial, and so the homotopy is reduction decidable. $\square$

Let $S_\emptyset$ be the empty set. Then any homotopy data triple $(\alpha, \tau, S_\emptyset)$ gives a homotopy where the third homotopy move is disallowed. Thus the homotopy relation is generated by isomorphism and the first two homotopy moves.

The homotopy relation induces an equivalence relation on the set of isomorphism classes of $\mathcal{P}(\alpha)$. We call a first homotopy move or second homotopy move *reducing* if it removes letters from a nanophrase. For two nanophrases p and q we write $p \geq q$ if either p is equal to q or q is derivable from p by a sequence of reducing homotopy moves. The following lemma shows that for the homotopy given by $(\alpha, \tau, S_\emptyset)$, the set of isomorphism classes satisfies the Confluence condition of Newman's Diamond Lemma (Lemma 5.13).

Lemma 6.5. *Let p be a nanophrase in $\mathcal{P}(\alpha)$. Suppose p_1 and p_2 are the results of applying different reducing moves to p under the homotopy given by $(\alpha, \tau, S_\emptyset)$. Then there exists a nanophrase p' such that $p_1 \geq p'$ and $p_2 \geq p'$.*

Proof. If the sets of letters involved in the two moves do not intersect, then it is clear that the moves can be applied in either order and give the same result in both cases. So we only need to check the cases where the sets of letters do intersect.

If both moves are first homotopy moves, the set of letters can not intersect.

If one move is a first homotopy move and the other a second homotopy move, then we must have the case $xABBAy$ where $|B| = \tau(|A|)$. Applying the second homotopy move gives xy . On the other hand, applying the first homotopy move gives $xAAy$. We can then apply another first homotopy move to get xy which was the result of applying the second homotopy move. Thus in this case the Confluence condition holds.

If both moves are second homotopy moves, then we must have the case $xABCyCBz$, where $|A| = \tau(|B|) = |C|$. Then one of the moves removes A and B to give $xCyCz$. The other move removes B and C to give $xAyAz$. However, these nanophrases are isomorphic because $|A|$ and $|C|$ are equal. Thus in this case the Confluence condition also holds. $\square$

As the rank of a nanophrase is finite and decreases under a reducing move, the Finiteness condition of Newman's Diamond Lemma is also satisfied. Thus, by Newman's Diamond Lemma (Lemma 5.13), the homotopy given by $(\alpha, \tau, S_\emptyset)$ has a normal form (up to isomorphism). Given a nanophrase p in $\mathcal{P}(\alpha)$, the normal form is calculated by applying reducing homotopy moves wherever they appear. When no more reducing homotopy moves are applicable, the normal form has been reached. In particular, it is clear that the normal form is minimizing.

We remark that Manturov noted the existence of this normal form in the specific case where α contains a single element in [10].

By Proposition 6.4, we have the following proposition.

Proposition 6.6. *Let $(\alpha, \tau, S_\emptyset)$ be a homotopy data triple. The homotopy that it gives is reduction decidable and equality decidable.*

In particular, there are exactly two prime data triples for which S is $S_\emptyset$. The first is $(\{a\}, \tau_{id}, S_\emptyset)$ where τ_{id} is the identity map. The second is $(\{a, b\}, \tau_a, S_\emptyset)$ where τ_a maps a to b (and therefore maps b to a).

Remark 6.7. We note that Theorem 3.8 of [9] would imply the existence of a minimizing normal form for the homotopy given by (α_F, τ_F, S_F) (defined in Example 4.3). However, we found a counter-example to the theorem which we explained in [7].

7. DETECTING IRREDUCIBILITY

In this section we look at ways of determining irreducibility of components of nanophrases.

7.1. Component length. In [5], we noted that the number of letters in a component modulo 2 is a homotopy invariant of Gauss phrases. Fukunaga made the same observation for nanophrases under homotopies with diagonal S in [4]. In fact, the number of letters in a component modulo 2 is a homotopy invariants of nanophrases for any choice of S .

Therefore, if the i th component of a nanophrase p contains an odd number of letters it cannot be i -reducible.

Example 7.1. Let p be the nanophrase $ABC|A|B|C$. Each component of p has an odd number of letters. Then, irrespective of the projections of the letters A , B and C , p is irreducible under any nanophrase homotopy.

7.2. Linking matrix. In [5], we defined a homotopy invariant of Gauss phrases called the *linking matrix*. In [2], Fukunaga defined the *linking vector* of a nanophrase and proved that it is invariant under homotopies with diagonal S . As we observed in [5], our linking matrix is equivalent to the linking vector of a Gauss phrase.

Using our linking matrix terminology, we now recall Fukunaga's linking vector invariant. We observe that it is invariant for any choice of S in the homotopy triple, not just for diagonal S .

We fix a homotopy data triple (α, τ, S) . Let π be the multiplicative abelian group generated by elements of α such that $a\tau(a) = 1$ for all a in α . This group was defined by Turaev in [15].

The *linking matrix* $L(p)$ of an n -component nanophrase is a symmetric $n \times n$ matrix with elements in π . The elements of $L(p)$ are denoted by $l_p(i, j)$. We define $l_p(i, i)$ to be 1 for all i and define $l_p(i, j)$ by

$$l_p(i, j) = \prod_{X \in \mathcal{A}_{ij}} |X|$$

where $\mathcal{A}_{ij}$ denotes the subset of letters appearing in p which appear both in the i th and j th components of p . Note that by definition, $l_p(i, j)$ is equal to $l_p(j, i)$ for all i and j . Fukunaga's linking vector is the vector $(l_p(1, 2), l_p(1, 3), \dots, l_p(1, n), l_p(2, 3), \dots, l_p(n-1, n))$. In the case where S is diagonal, Fukunaga showed in [2] that the linking matrix $L(p)$ is invariant. This is easily extended to the general case.

Proposition 7.2. *The linking matrix $L(p)$ is a homotopy invariant of p .*

Proof. After noting that the sets $\mathcal{A}_{ij}$ do not change under the third homotopy move for any S , the proof is the same as in [2]. $\square$

Example 7.3. Let p be the nanophrase $ABC|AC|B$ where $|A|$ is a , $|B|$ is b and $|C|$ is c . Then $L(p)$ is given by

$$\begin{pmatrix} 1 & ac & b \\ ac & 1 & 1 \\ b & 1 & 1 \end{pmatrix}.$$

Let $L_i(p)$ be the i th row of $L(p)$. We call $L_i(p)$ the *linking vector* of the i th component of p . Note that if the i th component of p is empty, then every element of $L_i(p)$ will be 1. Thus if there is an element in $L_i(p)$ which is not 1, then we can conclude that p is not i -reducible.

Example 7.4. Consider again the nanophrase p given in Example 7.3. From the linking matrix of p we can see that both the first and third rows contain the element b which is not 1 in π . Thus p is not 1-reducible or 3-reducible. Furthermore, if a is not equal to $\tau(c)$ then ac is not 1 in π and p is not 2-reducible either.

7.3. Subphrase invariants. Let p be an n -component nanophrase and let i be an element of $\hat{n}$. Let O be some subset of $\hat{n} - \{i\}$. Then, by Lemma 3.7, $x(p, O)$ is an invariant of p . As i is not in O , $x(p, O)$ contains a component corresponding to the i th component of p . Let j be the index of this component in $x(p, O)$. Then if $x(p, O)$ is not j -reducible, p is not i -reducible.

In particular, when O is the set $\hat{n} - \{i\}$, then $x(p, O)$ is a nanoword which we denote $w_i(p)$. If $w_i(p)$ is not contractible, p cannot be i -reducible.

Example 7.5. Let p be a nanophrase over α_1 given by $ABACDECBD|EF$. Then $w_1(p)$ is $ABACDCBD$ and $w_2(p)$ is $\emptyset$.

We consider p under the homotopy given by the homotopy data triple (α_G, τ_G, S_G) given in Example 4.3. This homotopy is the open Gauss word homotopy. By Example 6.2 of [6] we know that $ABACDCBD$ is not contractible. Therefore p is not 1-reducible.

7.4. V invariant. We defined the S_o invariant for Gauss phrases in [5]. Here we generalize the invariant to nanophrases for arbitrary α , τ and S . We rename the invariant V in order to avoid confusion with Fukunaga's generalization of the S_o invariant [3].

We write K_n for $(\mathbb{Z}/2\mathbb{Z})^n$. Let p be an n -component nanophrase and let i be an element of $\hat{n}$. Let $\mathcal{A}_{ii}(p)$ be the subset of letters appearing in p such that both occurrences of the letter appear in the i th component of p . Let A be a letter in $\mathcal{A}_{ii}(p)$. Then the i th component of p has the form $xAyAz$ for some, possibly empty, words x , y and z . We define the *linking vector* of A , written $\overrightarrow{l(A)}$, to be a vector $\vec{v}$ in K_n . The j th element of $\vec{v}$ is, modulo 2, the number of letters that appear once in y and for which the other occurrence of the letter appears in the j th component of p .

Let $\mathcal{A}_{i,a}$ be the subset of $\mathcal{A}_{ii}(p)$ consisting of letters A such that $|A|$ is equal to a . For each element a in α we define a map $d_{i,a}$ from $K_n - \{\vec{0}\}$ to $\mathbb{Z}$, given by

$$d_{i,a}(\vec{v}) = \sharp\{X \in \mathcal{A}_{i,a} \mid \overrightarrow{l(X)} = \vec{v}\},$$

where $\sharp$ means the number of elements in the set.

Proposition 7.6. *If $\tau(a)$ is equal to a then $d_{i,a}(\vec{v})$ modulo 2 is a homotopy invariant of p . If $\tau(a)$ is not equal to a then $d_{i,a}(\vec{v}) - d_{i,\tau(a)}(\vec{v})$ is a homotopy invariant of p .*

Proof. The proof is similar to that of the invariance of S_o for Gauss phrases in [5].

The point is that under a homotopy move, the linking vector of any letter in $\mathcal{A}_{ii}(p)$ does not change and so is counted the same in $d_{i,a}$, unless the letter is added or removed by the move.

If the letter is added or removed by the first homotopy move then the linking vector of the letter is $\vec{0}$. As $\vec{0}$ is not in the domain of $d_{i,a}$, $d_{i,a}$ is unchanged.

Under the second homotopy move two letters are added or removed. If one of the letters is in $\mathcal{A}_{ii}(p)$ then both are. In this case, both letters have the same linking vector $\vec{v}$. If the projection of neither letter is a then $d_{i,a}$ is unchanged. If not, the projection of one letter is a and the other is $\tau(a)$. If a is equal to $\tau(a)$, then we count the vector twice in $d_{i,a}$ and so, modulo 2, $d_{i,a}$ is unchanged. If a is not equal to $\tau(a)$, then one letter contributes to $d_{i,a}$ and the other letter contributes to $d_{i,\tau(a)}$. Thus $d_{i,a}(\vec{v}) - d_{i,\tau(a)}(\vec{v})$ is unchanged by the move. $\square$

We define $V^{i,a}(\vec{v})$ to be $d_{i,a}(\vec{v})$ modulo 2 if a is equal to $\tau(a)$ and $d_{i,a}(\vec{v}) - d_{i,\tau(a)}(\vec{v})$ otherwise. We fix a subset α_0 of α which contains exactly one element from each orbit of τ . Turaev calls this an *orientation* of α . Now note that if $\tau(a)$ is not equal to a then $V^{i,\tau(a)}(\vec{v})$ is equal to $-V^{i,a}(\vec{v})$ for all $\vec{v}$ in $K_n - \{\vec{0}\}$. Thus V is completely defined by the maps $V^{i,a}$ for a in α_0 .

Example 7.7. Let α be the set $\{a, b, c\}$. Let τ map a to b and c to itself. Let S be the set $\{(a, b, c), (c, b, a)\}$.

Let p be a nanophrase over α given by $ADBAEBFG|CDCF|EG$ where $|A| = |D| = a$, $|B| = |F| = b$ and $|C| = |E| = |G| = c$. Note that the linking matrix of p is trivial. Thus, using the linking matrix, p is indistinguishable from the trivial 3-component nanophrase $\emptyset|\emptyset|\emptyset$.

Now $\mathcal{A}_{11}(p)$ is the set $\{A, B\}$, $\mathcal{A}_{22}(p)$ is the set $\{C\}$ and $\mathcal{A}_{33}(p)$ is empty. Calculating the linking vectors of A , B and C , we see that $\overrightarrow{l(A)}$ is $(1, 1, 0)$, $\overrightarrow{l(B)}$ is $(1, 0, 1)$. and $\overrightarrow{l(C)}$ is $(1, 0, 0)$. Then, $d_{1,a}(\vec{v})$ is equal to 1 if $\vec{v}$ is equal to $(1, 1, 0)$ and 0 otherwise. Similarly, $d_{1,b}(\vec{v})$ is equal to 1 if $\vec{v}$ is equal to $(1, 0, 1)$ and 0 otherwise. Furthermore, $d_{2,c}(\vec{v})$ is equal to 1 if $\vec{v}$ is equal to $(1, 0, 0)$ and 0 otherwise. All the other maps $d_{i,x}$ map every vector to 0.

Therefore $V^{1,a}(\vec{v})$ is equal to 1 if $\vec{v}$ is equal to $(1, 1, 0)$, -1 if $\vec{v}$ is equal to $(1, 0, 1)$ and 0 otherwise. As b is equal to $\tau(a)$, $V^{1,b}(\vec{v})$ is equal to $-V^{1,a}(\vec{v})$. Thus $V^{1,b}(\vec{v})$ is equal to -1 if $\vec{v}$ is equal to $(1, 1, 0)$, 1 if $\vec{v}$ is equal to $(1, 0, 1)$ and 0 otherwise. Also, $V^{2,c}(\vec{v})$ is equal to 1 if $\vec{v}$ is equal to $(1, 0, 0)$ and 0 otherwise. For all other pairs (i, x) , $V^{i,x}(\vec{v})$ is 0 for all $\vec{v}$.

Since $\emptyset|\emptyset|\emptyset$ has trivial maps for all $V^{i,x}$, we may conclude that p and $\emptyset|\emptyset|\emptyset$ are not homotopic.

Remark 7.8. In the case of Gauss phrases, α contains a single element a and τ is the identity map. Thus for the i th component we just have a single map $V^{i,a}$ from $K_n - \{\vec{0}\}$ to $\mathbb{Z}/2\mathbb{Z}$. In [5] we defined $B_i(p)$ to be the preimage of 1 under the map $V^{i,a}$ and then defined S_o to be an n -tuple where the i th element is $B_i(p)$.

Remark 7.9. In [3] Fukunaga generalized the Gauss phrase S_o invariant to any homotopy with diagonal S . His invariant is also called S_o . However, his invariant is equivalent to the invariant we call U which is defined in Section 7.5.

If, for a given p , $\mathcal{A}_{ii}(p)$ is empty then $V^{i,a}(\vec{v})$ is 0 for all vectors $\vec{v}$. Thus p is not i -reducible if there exists some a in α and some vector $\vec{v}$ such that $V^{i,a}(\vec{v})$ is non-zero.

Example 7.10. Consider again the nanophrase p defined in Example 7.7. From our calculations in that example we know that $V^{1,a}((1, 1, 0))$ is non-zero. We also know that $V^{2,c}((1, 0, 0))$ is non-zero. Therefore, p is neither 1-reducible nor 2-reducible.

Note that since the linking matrix of p is trivial, we could not determine this information from the linking matrix.

7.5. U invariant. In [15] Turaev defined the self-linking function of a nanoword, an invariant for nanowords under homotopies with diagonal S . This invariant is a generalization of the u -polynomial defined by Turaev for flat virtual knots (also known as virtual strings) [13]. Based on the constructions of the self-linking function for nanowords and the invariant V we define a new invariant for nanophrases under homotopies with diagonal S which we call U .

Let π be the group defined in Section 7.2. Let p be an n -component nanophrase and i be an integer in $\hat{n}$. Let A and B be two letters in $\mathcal{A}_{ii}(p)$. We define $n(A, B)$ to be 1 if the i th component of p has the form $\dots A \dots B \dots A \dots B \dots$, $n(A, B)$ to be -1 if the i th component of p has the form $\dots B \dots A \dots B \dots A \dots$ and $n(A, B)$ to be 0 otherwise. In particular $n(A, A)$ is defined to be 0 for all A in $\mathcal{A}_{ii}(p)$.

For A in $\mathcal{A}_{ii}(p)$, we define $\overrightarrow{l_u(A)}$ to be a vector in π^n . The i th element of $\overrightarrow{l_u(A)}$ is defined to be

$$\prod_{B \in \mathcal{A}_{ii}(p)} |B|^{n(A, B)}.$$

Writing the i th component of p as $xAyAz$ for some, possibly empty, words x , y and z , we define the j th element of $\overrightarrow{l_u(A)}$ (j not equal to i) to be

$$\prod_{B \in \mathcal{A}_{ij}(p), B \in y} |B|.$$

Then for each a in α and for each i , let $\mathcal{A}_{i,a}$ be the subset of $\mathcal{A}_{ii}(p)$ consisting of letters A such that $|A|$ is equal to a . Let $\vec{1}$ be the trivial vector in π^n for which

every element is 1. For each element a in α we define a map $e_{i,a}$ from $\pi^n - \{\vec{1}\}$ to $\mathbb{Z}$, given by

$$e_{i,a}(\vec{v}) = \sharp\{X \in \mathcal{A}_{i,a} \mid \overrightarrow{l_u(X)} = \vec{v}\},$$

where $\sharp$ means the number of elements in the set.

Remark 7.11. When n is 1, p is a nanoword. Then the self-linking class $[a]_p$ defined in [15] can be derived from $e_{1,a}$ by

$$[a]_p = \sum_{\vec{v} \in \pi - \{\vec{1}\}} e_{1,a}(\vec{v}) \vec{v}.$$

In light of Remark 7.11, the following proposition is a generalization of Theorem 6.1.1 in [15].

Proposition 7.12. *If $\tau(a)$ is equal to a then $e_{i,a}$ modulo 2 is a homotopy invariant of p . If $\tau(a)$ is not equal to a then $e_{i,a} - e_{i,\tau(a)}$ is a homotopy invariant of p .*

Proof. It is sufficient to prove invariance under a single isomorphism or homotopy move.

It is clear that $\overrightarrow{l_u(A)}$ does not change under isomorphism. Thus $e_{i,a}$ is unchanged under isomorphism.

We now check the first homotopy move. Suppose A is the letter removed under the move. Then for any other letter B , $\overrightarrow{l_u(B)}$ is unchanged by the removal of A . On the other hand, $\overrightarrow{l_u(A)}$ is the trivial vector $\vec{1}$ and so $\overrightarrow{l_u(A)}$ cannot be in the domain of $e_{i,a}$. Thus $e_{i,a}$ does not change under the first homotopy move.

We now check the second homotopy move. Suppose A and B are the letters removed under the move. Let C be some other letter and consider how $\overrightarrow{l_u(C)}$ is affected by the move. We note that if A contributes to $\overrightarrow{l_u(C)}$ then so does B . Since $|A| \cdot |B|$ equals 1, if A and B contribute to $\overrightarrow{l_u(C)}$, their contributions cancel. Thus $\overrightarrow{l_u(C)}$ is unchanged by the removal of A and B . On the other hand, A is in $\mathcal{A}_{ii}(p)$ if and only if B is. In the case where they are both in $\mathcal{A}_{ii}(p)$, $\overrightarrow{l_u(A)}$ is equal to $\overrightarrow{l_u(B)}$. If neither A nor B project to a , $e_{i,a}$ is unchanged under the move. Now if the projection of one of A or B is a , then the projection of the other is $\tau(a)$. Without loss of generality, we may assume that $|A|$ is a . If a is equal to $\tau(a)$ then both A and B contribute to $e_{i,a}(\overrightarrow{l_u(A)})$ and, modulo 2, the contributions cancel. If a is not equal to $\tau(a)$ then A contributes to $e_{i,a}(\overrightarrow{l_u(A)})$ and B contributes to $e_{i,\tau(a)}(\overrightarrow{l_u(A)})$. Thus $e_{i,a}(\overrightarrow{l_u(A)}) - e_{i,\tau(a)}(\overrightarrow{l_u(A)})$ is unchanged by the move.

We now check the third homotopy move. Suppose A , B and C are the letters involved in the move and before the move we have the pattern $xAByACzBCt$. For any other letter D , if the letters A , B or C do not contribute to $\overrightarrow{l_u(D)}$, then $\overrightarrow{l_u(D)}$ is unaffected by the move. If the letters A , B or C do contribute to $\overrightarrow{l_u(D)}$, then, as π is abelian, the order of the contributing letters is irrelevant and so $\overrightarrow{l_u(D)}$ is unchanged by the move.

Now consider A . Before the move B contributes to $\overrightarrow{l_u(A)}$ and C does not. After the move C contributes to $\overrightarrow{l_u(A)}$ and B does not. However, $|B|$ is equal to $|C|$ and so the contributions of B and C to $\overrightarrow{l_u(A)}$ are equal. Thus $\overrightarrow{l_u(A)}$ does not change under the move.

Now consider B . Before the move, both A and C contribute to $\overrightarrow{l_u(B)}$ but their contributions cancel. After the move, neither A nor C contribute to $\overrightarrow{l_u(B)}$. Thus $\overrightarrow{l_u(B)}$ does not change under the move.

The case of C is symmetric to the case of A and so $\overrightarrow{l_u(C)}$ does not change under the move.

Thus $e_{i,a}$ is invariant under the third homotopy move. $\square$

We define $U^{i,a}(\vec{v})$ to be $e_{i,a}(\vec{v})$ modulo 2 if a is equal to $\tau(a)$ and $e_{i,a}(\vec{v}) - e_{i,\tau(a)}(\vec{v})$ otherwise. As we did for the V invariant, we pick an orientation α_0 of α (α_0 contains exactly one element from each orbit of α under τ). Now note that if $\tau(a)$ is not equal to a then $U^{i,\tau(a)}(\vec{v})$ is equal to $-U^{i,a}(\vec{v})$ for all $\vec{v}$ in $\pi^n - \{\vec{1}\}$. Thus U is completely determined by the maps $U^{i,a}$ for a in α_0 .

Example 7.13. Let α be the set $\{a, b, c, d\}$. Let τ map a to b and c to d . Let S be the diagonal of α .

Let p be a nanophrase over α given by $ACDEABFB|CE|DF$ where $|A| = |C| = |E| = a$, $|B| = c$, $|D| = |F| = d$. Let q be the nanophrase $CDEF|CE|DF$ where the projections are the same as for p . Then it is easy to check that the linking matrix cannot distinguish p and q .

We calculate U for p . We see that $\overrightarrow{l_u(A)}$ is $(1, a^2, c^{-1})$ and $\overrightarrow{l_u(B)}$ is $(1, 1, c^{-1})$. Thus $U^{1,a}(\vec{v})$ is 1 if $\vec{v}$ is $(1, a^2, c^{-1})$ and 0 otherwise. Also, $U^{1,c}(\vec{v})$ is -1 if $\vec{v}$ is $(1, 1, c^{-1})$ and 0 otherwise. For all other pairs (i, x) , $U^{i,x}$ is 0 for all vectors.

On the other hand, for q , the maps $U^{i,x}$ are trivial for all i and x . Thus p and q are not homotopic.

Remark 7.14. Having written this section we discovered that Fukunaga had also independently generalized the Gauss phrase S_o invariant to any homotopy with diagonal S in [3]. His invariant is called S_o .

The initial definition of Fukunaga's S_o invariant, appearing in an early version of [3], was weaker than our U invariant. However, we realised that by a slight modification of the definition, Fukunaga's S_o could be strengthened. This modified definition is now the one that appears in [3]. In Section 7.6 we will show that our U invariant is equivalent to Fukunaga's S_o invariant.

For any nanophrase we can calculate $V^{i,a}$ from $U^{i,a}$. To do so, we define a map x from π to $\mathbb{Z}/2\mathbb{Z}$ as follows. Any element g of π can be written uniquely in the form

$$(7.1) \quad \prod_{a \in \alpha_0} a^{i_a}$$

where i_a is an integer depending on a . Then we define $x(g)$ to be the sum of the exponents in Equation (7.1) modulo 2, that is

$$x(g) = \sum_{a \in \alpha_0} i_a \pmod{2}.$$

The map x then induces a map from π^n to K^n , which we also call x , by applying x elementwise. For a vector $\vec{v}$ in K^n , let I_v be the preimage of $\vec{v}$ under x . Then

$$d_{i,a}(\vec{v}) = \sum_{\vec{u} \in I_v} e_{i,a}(\vec{u}).$$

Thus if a is equal to $\tau(a)$,

$$V^{i,a}(\vec{v}) = \sum_{\vec{u} \in I_v} U^{i,a}(\vec{u}) \pmod{2},$$

and if a is not equal to $\tau(a)$,

$$V^{i,a}(\vec{v}) = \sum_{\vec{u} \in I_v} U^{i,a}(\vec{u}).$$

We have already noted in Example 4.3 that there are only two prime homotopy data triples for which S is diagonal. For (α_G, τ_G, S_G) (using the notation of Example 4.3), the V invariant and U invariant are equivalent. For (α_F, τ_F, S_F) , the U invariant is stronger than the V invariant.

Example 7.15. Let p be the nanophrase $ABCA|BC$ where $|A|$, $|B|$ and $|C|$ are all a an element in α_F .

We calculate the V invariant for p . As the linking vector for A , $\overrightarrow{l(A)}$ is $(0, 0)$, and A is the only letter in p for which both occurrences appear in the same component, we conclude that $d_{i,a}(\vec{v})$ is 0 for i equal to 1 or 2 and for all $\vec{v}$ in $K_n - \{\vec{0}\}$. So $V^{i,a}(\vec{v})$ is 0 for i equal to 1 or 2 and for all $\vec{v}$ in $K_n - \{\vec{0}\}$. Thus, using the V invariant, p is indistinguishable from the nanophrase $\emptyset|\emptyset$.

We now calculate the U invariant for p . In this case, the linking vector for A , $\overrightarrow{l_u(A)}$ is $(1, a^2)$. Note that as a is not equal to $\tau(a)$, a^2 is not equal to 1, so $(1, a^2)$ is a non-trivial vector in π^2 . Thus $e_{1,a}((1, a^2))$ is equal to 1 and so $U^{1,a}((1, a^2))$ is equal to 1. Therefore p is not homotopic to $\emptyset|\emptyset$.

For completeness we note that for all other vectors $\vec{v}$ in $\pi^n - \{\vec{1}\}$, $U^{1,a}(\vec{v})$ is equal to 0. For all vectors $\vec{v}$ in $\pi^n - \{\vec{1}\}$, $U^{2,a}(\vec{v})$ is equal to 0.

For a given p , if $\mathcal{A}_{ii}(p)$ is empty then for all $\vec{v}$, $U^{i,a}(\vec{v})$ is 0. Thus p cannot be i -reducible if there exists some a in α and some vector $\vec{v}$ for which $U^{i,a}(\vec{v})$ is non-zero. Indeed, the U invariant can sometimes detect i -irreducibility where the V invariant cannot.

Let p be a nanophrase and let w be $w_i(p)$ for some i , where $w_i(p)$ is the nanoword derived from the i th component of p which was defined in Section 7.3. For any g in π let I_g be the set

$$\{\vec{v} \in \pi^n - \{\vec{1}\} \mid [\vec{v}]_i = g\}$$

where $[\vec{v}]_i$ denotes the i th element of $\vec{v}$. Then it is easy to check that $U^{1,a}(g)$ for w is given by

$$\sum_{\vec{v} \in I_g} U^{i,a}(\vec{v})$$

where the sum is taken modulo 2 if a is equal to $\tau(a)$.

Recall that any element g of π can be written in the form given in (7.1). We define $\gamma_a(g)$ to be the exponent of a when g is written in this form (that is, $\gamma_a(g)$ is equal to i_a in (7.1)). For a nanophrase, we define $\delta_a(U^{i,b})$ by

$$(7.2) \quad \delta_a(U^{i,b}) = \sum_{g \in \pi - \{1\}} \sum_{\vec{v} \in I_g} U^{i,b}(\vec{v}) \gamma_a(g)$$

where the sum is taken modulo 2 if a is equal to $\tau(a)$. In [15], Turaev gave necessary and sufficient conditions for a self-linking function to be realizable as a self-linking function of a nanoword. These conditions translate to the condition that

$$\delta_a(U^{1,b}) + \delta_b(U^{1,a}) = 0$$

for all a and b in α_0 and

$$\delta_a(U^{1,a}) = 0$$

for all a in α_0 .

This gives some necessary conditions for a set of maps $U^{i,a}$ to be realizable as the U invariant of a nanophrase. In fact these conditions are also sufficient.

Proposition 7.16. *For each a in α_0 , let $U^{i,a}$ be a map from $\pi^n - \{\vec{1}\}$ to $\mathbb{Z}$, if a is not equal to $\tau(a)$, and to $\mathbb{Z}/2\mathbb{Z}$ otherwise. If, for each i , the maps defined in (7.2) satisfy*

$$\delta_a(U^{i,b}) + \delta_b(U^{i,a}) = 0$$

for all a and b in α_0 and

$$\delta_a(U^{i,a}) = 0$$

for all a in α_0 , then the maps $U^{i,a}$ are the U invariant of some nanophrase.

Proof. This can be proved by combining the arguments in Theorem 6.3.1 of [15] and Proposition 5.5 and Proposition 5.7 of [5]. We omit the details. $\square$

Remark 7.17. By a similar argument to that given in Proposition 5.7 of [5], the linking matrix and U invariants can be shown to be independent.

7.6. U invariant and Fukunaga's S_o invariant. In [3], Fukunaga generalized our S_o invariant for Gauss phrases defined in [5]. His invariant is also called S_o . In this subsection we recall the definition of Fukunaga's S_o invariant and show that it is equivalent to our U invariant.

We fix a homotopy (α, τ, S) with diagonal S . An orbit of τ is called a *free orbit* if the orbit contains two elements of α . Otherwise, the orbit contains only one element and is called a *fixed orbit*. Let l be the number of free orbits of τ and m be the number of fixed orbits of τ . We denote the orbits of τ by $\widehat{a}_i$ where, for i running from 1 to l , $\widehat{a}_i$ is a free orbit of τ and for i running from $l+1$ to m , $\widehat{a}_i$ is a fixed orbit. For each orbit $\widehat{a}_i$ of τ we fix a representative element which we denote a_i . The set of representative elements is an orientation of α which we denote α_0 .

Let p be an n -component nanophrase. For a letter A in p , $\varepsilon(A)$ is defined as follows:

$$\varepsilon(A) = \begin{cases} 1 & \text{if } |A| = a_j \text{ and } 1 \leq j \leq l+m \\ -1 & \text{if } |A| = \tau(a_j) \text{ and } 1 \leq j \leq l. \end{cases}$$

Let $K_{i,j}$ be $\mathbb{Z}$ if i and j are both less or equal to l and let $K_{i,j}$ be $\mathbb{Z}/2\mathbb{Z}$ otherwise. Let K be $K_{1,1} \times K_{1,2} \times \dots \times K_{1,l+m} \times K_{2,1} \times \dots \times K_{l+m,l+m}$. Elements of K are considered to be row vectors. Let $\vec{r}_{s,t}$ denote the element of K which has a 1 in the column corresponding to $K_{s,t}$ and 0 in every other column. Let $\vec{0}$ denote the zero vector in K .

For a nanophrase p , let $\mathcal{A}(p)$ be the α -alphabet associated with p . We now extend the definition of $n(A, B)$ given in Section 7.5 to any two letters A and B in $\mathcal{A}(p)$. We define $n(A, B)$ to be 1 if p has the form $\dots A \dots B \dots A \dots B \dots$, $n(A, B)$ to be -1 if p has the form $\dots B \dots A \dots B \dots A \dots$ and $n(A, B)$ to be 0 otherwise. As before, $n(A, A)$ is defined to be 0 for all A in $\mathcal{A}(p)$.

For any letter B in $\mathcal{A}(p)$, let $I_1(B)$ denote the index of the component in which the first occurrence of B appears. Similarly, let $I_2(B)$ denote the index of the component in which the second occurrence of B appears.

For any letter A in $\mathcal{A}(p)$, Fukunaga defines $\sigma_j(A, B)$ by

$$\sigma_j(A, B) = \begin{cases} \vec{r}_{s,t} & \text{if } |A| \in \widehat{a}_s, |B| = a_t, n(A, B) = 1 \text{ and } I_2(B) = j, \\ \vec{r}_{s,t} & \text{if } |A| \in \widehat{a}_s, |B| = \tau(a_t), n(A, B) = -1 \text{ and } I_1(B) = j, \\ -\vec{r}_{s,t} & \text{if } |A| \in \widehat{a}_s, |B| = \tau(a_t), n(A, B) = 1 \text{ and } I_2(B) = j, \\ -\vec{r}_{s,t} & \text{if } |A| \in \widehat{a}_s, |B| = a_t, n(A, B) = -1 \text{ and } I_1(B) = j, \\ \vec{0} & \text{otherwise.} \end{cases}$$

Now, for any letter A in $\mathcal{A}(p)$, Fukunaga defines $l_j(A)$ by

$$l_j(A) = \sum_{X \in \mathcal{A}(p)} \sigma_j(A, X)$$

and $l(A)$ as an n -tuple given by

$$l(A) = (l_1(A), l_2(A), \dots, l_n(A)).$$

Note that $l(A)$ is in K^n .

For a vector $\vec{v}$ in K^n , let $v_{i,s,t}$ be the value of the column corresponding to $K_{s,t}$ in the i th component of $\vec{v}$. Let $\vec{0}_n$ be the zero vector in K^n . Now, for any vector $\vec{v}$ in $K^n - \vec{0}_n$, we define the *type* of $\vec{v}$ as follows. We say $\vec{v}$ is of type (i) if $v_{i,r,s}$ being non-zero implies r is less than or equal to l , for all i, r and s . We say $\vec{v}$ is of type (ii) if $v_{i,r,s}$ being non-zero implies r is greater than l , for all i, r and s . Otherwise, we say $\vec{v}$ is of type (iii).

For an integer i and a vector $\vec{v}$ in K^n we define $\eta(i, \vec{v})$ by

$$\eta(i, \vec{v}) = \sum_{A \in \mathcal{A}_{ii}(p), l(A)=\vec{v}} \varepsilon(A)$$

Then Fukunaga defines a map B_i from $K^n - \{\vec{0}_n\}$ to $\mathbb{Z}$ or $\mathbb{Z}/2\mathbb{Z}$, depending on the type of vector, as follows:

$$B_i(\vec{v}) = \begin{cases} \eta(i, \vec{v}) & \text{if } \vec{v} \text{ is of type (i),} \\ \eta(i, \vec{v}) \mod 2 & \text{if } \vec{v} \text{ is of type (ii),} \\ 0 & \text{otherwise.} \end{cases}$$

Finally, $S_o(p)$ is the n -tuple of maps given by

$$S_o(p) = (B_1, B_2, \dots, B_n).$$

In [3], Fukunaga proved that $S_o(p)$ is a homotopy invariant of p .

In order to show that Fukunaga's S_o invariant and the U invariant are equivalent we need some preparation.

For any element g in π , where π is the group defined in Section 7.2, we can uniquely write g in the form

$$g = \prod_{i=1}^{l+m} a_i^{c_i}$$

where c_i is in $\mathbb{Z}$ if i is less than or equal to l and c_i is in $\mathbb{Z}/2\mathbb{Z}$ otherwise. For each i running from 1 to l we define a map f_i from π to $\mathbb{Z}$. For each i running from $l+1$ to $l+m$ we define a map f_i from π to $\mathbb{Z}/2\mathbb{Z}$. In either case, $f_i(g)$ is defined to be c_i , the exponent of a_i when g is written in the form above.

Now, for an element a in α_0 we define a set of maps $h_{a,i,j}$ from π to K . For g , an element of π , $h_{a,i,j}$ maps g to $\vec{u}$ where the component of $\vec{u}$ corresponding to $K_{s,t}$ is given by

$$u_{s,t} = \begin{cases} 0 & \text{if } a \notin \widehat{a}_s, \\ f_t(g) & \text{if } a \in \widehat{a}_s \text{ and } j \leq i, \\ -f_t(g) & \text{if } a \in \widehat{a}_s \text{ and } i < j. \end{cases}$$

Next we define a set of maps $h_{a,i}$ from π^n to K^n . The map $h_{a,i}$ takes $(g_1, g_2, \dots, g_n)$ to $(h_{a,i,1}(g_1), h_{a,i,2}(g_2), \dots, h_{a,i,n}(g_n))$.

For a nanophrase p and for some integer i , let A be a letter in $\mathcal{A}_{ii}(p)$. Then it follows from the definitions that Fukunaga's $l(A)$ is equal to $h_{a,i}(\overrightarrow{l_u(A)})$.

Suppose that $|A|$ is in $\widehat{a}_r$ for some r . Recall that for a vector $\vec{v}$ in K^n , $v_{j,s,t}$ is the value of the column corresponding to $K_{s,t}$ in the j th component of $\vec{v}$. Now observe that by definition, $l(A)_{j,s,t}$ is equal to zero if s is not equal to r . In other words, if $l(A)$ is non-zero, the orbit of τ which contains $|A|$ is implicitly recorded in $l(A)$.

The following lemma shows that we can calculate U from Fukunaga's invariant.

Lemma 7.18. *With the above notation, for any vector $\vec{v}$ in $\pi^n - \{\vec{1}\}$, we have*

$$U^{i,a}(\vec{v}) = B_i(h_{a,i}(\vec{v})).$$

Proof. When a is not equal to $\tau(a)$, we have

$$\begin{aligned} U^{i,a}(\vec{v}) &= e_o^{i,a}(\vec{v}) - e_o^{i,\tau(a)}(\vec{v}) \\ &= \#\{X \in \mathcal{A}_{i,a} \mid \overrightarrow{l_u(X)} = \vec{v}\} - \#\{X \in \mathcal{A}_{i,\tau(a)} \mid \overrightarrow{l_u(X)} = \vec{v}\} \\ &= \sum_{A \in \mathcal{A}_{ii}(p), l(A)=h_{a,i}(\vec{v})} \varepsilon(A) \\ &= B_i(h_{a,i}(\vec{v})). \end{aligned}$$

The equality before last holds because, by the observation given before the statement of the lemma, $l(A)$ can only equal $h_{a,i}(\vec{v})$ if $|A|$ is equal to a or $\tau(a)$. The last equality holds because if a is not equal to $\tau(a)$, and $|A|$ equals a , $l(A)$ must be a type (i) vector.

When a is equal to $\tau(a)$, we have a similar calculation:

$$\begin{aligned} U^{i,a}(\vec{v}) &= e_o^{i,a}(\vec{v}) \pmod{2} \\ &= \#\{X \in \mathcal{A}_{i,a} \mid \overrightarrow{l_u(X)} = \vec{v}\} \pmod{2} \\ &= \sum_{A \in \mathcal{A}_{ii}(p), l(A)=h_{a,i}(\vec{v})} \varepsilon(A) \pmod{2} \\ &= B_i(h_{a,i}(\vec{v})). \end{aligned}$$

The last equality holds because if a is equal to $\tau(a)$, and $|A|$ equals a , $l(A)$ must be a type (ii) vector. $\square$

Let $\mathcal{K}$ denote the subset of $K^n - \{\vec{0}_n\}$ given by

$$\mathcal{K} = \{\vec{v} \in K^n - \{\vec{0}_n\} \mid \exists r \in \mathbb{Z} \text{ such that } \forall j, s, t, v_{j,s,t} \neq 0 \Rightarrow s = r\}.$$

By the observation given before Lemma 7.18 it is clear that for any nanophrase p and any letter A in $\mathcal{A}_{ii}(p)$, either $l(A)$ is $\vec{0}_n$ or $l(A)$ is in $\mathcal{K}$.

We define maps κ_i from $\mathcal{K}$ to π_n . Let $\vec{v}$ be a vector in $\mathcal{K}$. By definition there exists an r such that $v_{j,s,t}$ is non-zero implies s equals r . Denote that r by $r(\vec{v})$. Now let $\vec{u}$ be $\kappa_i(\vec{v})$ for some integer i . Then u_j , the j th component of $\vec{u}$, is given by

$$u_j = \begin{cases} \prod_{a_t \in \alpha_0} a_t^{v_{j,r,t}} & \text{if } j \geq i, \\ \prod_{a_t \in \alpha_0} a_t^{-v_{j,r,t}} & \text{if } j < i. \end{cases}$$

The following lemma shows that we can calculate Fukunaga's invariant from U .

Lemma 7.19. *Given a vector $\vec{v}$ in $K^n - \{\vec{0}_n\}$, write a for $a_{r(\vec{v})}$. Then we have*

$$B_i(\vec{v}) = U^{i,a}(\kappa_i(\vec{v})).$$

Proof. If $\vec{v}$ is of type (i) then a is not equal to $\tau(a)$. In this case we have

$$\begin{aligned} B_i(\vec{v}) &= \sum_{A \in \mathcal{A}_{ii}(p), l(A)=\vec{v}} \varepsilon(A) \\ &= \#\{X \in \mathcal{A}_{i,a} \mid l(A) = \vec{v}\} - \#\{X \in \mathcal{A}_{i,\tau(a)} \mid l(A) = \vec{v}\} \\ &= \#\{X \in \mathcal{A}_{i,a} \mid \overrightarrow{l_u(A)} = \kappa_i(\vec{v})\} - \#\{X \in \mathcal{A}_{i,\tau(a)} \mid \overrightarrow{l_u(A)} = \kappa_i(\vec{v})\} \\ &= e_o^{i,a}(\kappa_i(\vec{v})) - e_o^{i,\tau(a)}(\kappa_i(\vec{v})) \\ &= U^{i,a}(\kappa_i(\vec{v})). \end{aligned}$$

If $\vec{v}$ is of type (ii) then a is equal to $\tau(a)$. In this case we have a similar calculation:

$$B_i(\vec{v}) = \sum_{A \in \mathcal{A}_{ii}(p), l(A)=\vec{v}} \varepsilon(A) \pmod{2}$$

$$\begin{aligned}
&= \#\{X \in \mathcal{A}_{i,a} \mid l(A) = \vec{v}\} \pmod{2} \\
&= \#\{X \in \mathcal{A}_{i,a} \mid \overrightarrow{l_u(A)} = \kappa_i(\vec{v})\} \pmod{2} \\
&= e_o^{i,a}(\kappa_i(\vec{v})) \pmod{2} \\
&= U^{i,a}(\kappa_i(\vec{v})).
\end{aligned}$$

□

Combining Lemma 7.18 and Lemma 7.19 proves the following proposition.

Proposition 7.20. *Fukunaga's S_o invariant is equivalent to the U invariant.*

Remark 7.21. In [2], Fukunaga defined the T invariant for nanophrases which is invariant under homotopies with diagonal S . In [3], Fukunaga showed that his generalized S_o invariant is strictly stronger than his T invariant. In particular, in Proposition 5.4 of [3], Fukunaga showed how to calculate the T invariant from the S_o invariant. As our U invariant is equivalent to Fukunaga's S_o invariant, U is strictly stronger than the T invariant.

8. NANOMULTIPHRASES

An n -phrase *multiphrase* on an alphabet $\mathcal{A}$ is a sequence of n phrases on $\mathcal{A}$. The concepts of *Gauss multiphrase* and *nanomultiphrase* are defined in an analogous way to those of Gauss phrase and nanophrase. When writing a multiphrase we use the symbol ‘||’ to separate phrases. For example $A|B||AC||D|B|CD$ is a 3-phrase Gauss multiphrase on $\{A, B, C, D\}$. Note that the multiphrases $A||B$ and $A|\emptyset|B$ are different, because an empty component in a phrase is always written $\emptyset$. So $A||B$ is a 2-phrase multiphrase where each phrase has a single component. The multiphrase $A|\emptyset|B$ is a 1-phrase multiphrase where the single phrase has 3 components.

The 0-component phrase $\emptyset_P$ may also appear as a phrase in a nanomultiphrase. There is a unique 0-phrase nanomultiphrase which is written $\emptyset_M$.

The only phrase of a 1-phrase nanomultiphrase is necessarily a nanophrase. Thus we can identify 1-phrase nanomultiphrase with nanophrases.

We define isomorphism of nanomultiphrases and homotopy moves on nanomultiphrases in an analogous way to nanophrases. Homotopy of nanomultiphrases is then the equivalence relation generated by isomorphisms and homotopy moves. We note that the number of phrases and the number of components in each phrase are invariant under homotopy.

Let p_1 be an n_1 -component phrase and p_2 be an n_2 -component phrase. Their *concatenation*, written $p_1|p_2$, is the $(n_1 + n_2)$ -component phrase consisting of the components of p_1 followed by the components of p_2 .

Let $\mathcal{M}(\alpha)$ be the set of nanomultiphrases over α . There exists a natural map μ from $\mathcal{M}(\alpha)$ to $\mathcal{P}(\alpha)$ where a nanomultiphrase m is mapped to a nanophrase by concatenating the phrases of m to make a single phrase. The following lemma is clear from the definitions.

Lemma 8.1. *Let m_1 and m_2 be nanomultiphrases. If $m_1 \sim m_2$, then $\mu(m_1) \sim \mu(m_2)$ as nanophrases.*

Let m be a nanomultiphrase. Then we define the i th component of m to be the i th component of $\mu(m)$. We define $\text{nc}(m)$ to be $\text{nc}(\mu(m))$, which is equal to the sum of the number of components appearing in each phrase of m . For example, $\text{nc}(A|B||AC||D|B|CD)$ is 6.

Recall that in Section 3 we defined the *concatenating map* χ from $\mathcal{P}(\alpha)$ to $\mathcal{N}(\alpha)$. We extend χ to be a map from $\mathcal{M}(\alpha)$ to $\mathcal{P}(\alpha)$ as follows. Let m be an n -phrase nanomultiphrase. Then $\chi(m)$ is the n -component nanophrase p where the i th

component of p is the word given by concatenating the components of the i th phrase of m . For example, $\chi(A|B||AC||D|B|CD)$ is $AB|AC|DBCD$.

We fix a homotopy data triple (α, τ, S) and let its prime factors be denoted by (α_i, τ_i, S_i) for i running from 1 to k for some k . Let $\mathcal{M}_A(\alpha)$ be the set of pairs (m, θ) , where m is a nanomultiphrase over α and θ is a map from $\widehat{\text{nc}(m)}$ to $\hat{k}$ such that, for all i , $\theta(i) = \theta(i+1)$ implies the i th and $(i+1)$ th components of m belong to different phrases of m . We extend the definition of isomorphism and homotopy moves of nanophrases to $\mathcal{M}_A(\alpha)$ by saying that $(m, \theta) \sim (m', \theta')$ if $p \sim p'$ and $\theta = \theta'$.

Let $\mathcal{M}_R(\alpha)$ be the subset of $\mathcal{M}_A(\alpha)$ consisting of pairs (m, θ) such that for every letter X appearing in the i th component of m , $|X|$ is in $\alpha_{\theta(i)}$, for all i . We define the equivalence relation $\sim_K$ on $\mathcal{M}_R(\alpha)$ as follows. We say $(m, \theta) \sim_K (m', \theta')$ if there exists a sequence of elements of $\mathcal{M}_R(\alpha)$, (m_i, θ_i) for i running from 0 to r for some r , such that m_0 is m , m_r is m' and, for each i , m_i is related to m_{i+1} by a single homotopy move or an isomorphism.

Reductions and augmentations are defined on $\mathcal{M}_R(\alpha)$ in an analogous way to how we defined them for $\mathcal{P}_R(\alpha)$. However, we do not allow concatenating reductions that would concatenate components coming from different phrases. So if the first or last component of a phrase is reducible, the reduction is made by a simple reduction, irrespective of θ . Let $\sim_M$ be the equivalence relation on $\mathcal{P}_M$ generated by $\sim_K$, reductions and augmentations.

We extend the map ψ to be a map from $\mathcal{P}(\alpha)$ to $\mathcal{M}_R(\alpha)$ as follows. The 0-component nanophrase $\emptyset_P$ is mapped to $\emptyset_M$. Any other element p of $\mathcal{P}(\alpha)$ is an n -component nanophrase in $\mathcal{P}(\alpha)$ with n greater than 0. We map each component w_i of p to a phrase. If w_i is $\emptyset$ then we map w_i to $\emptyset_P$. Otherwise, there is a unique integer $r > 0$, a unique sequence of words $x_1, x_2, \dots, x_r$ and a unique map θ_i from $\hat{r}$ to $\hat{k}$ such that

- (1) each x_j is not $\emptyset$;
- (2) w_i is equal to $x_1 x_2 \dots x_r$;
- (3) for all j , if X appears in w_j , $|X|$ is in $\alpha_{\theta_i(j)}$;
- (4) for any j from 1 to $r-1$, $\theta_i(j)$ is not equal to $\theta(j+1)$.

Then w_i maps to the phrase $x_1|x_2|\dots|x_r$. Write p_i for the image of w_i . Then p is mapped to $p_1|p_2|\dots|p_n$ which we label m . We define θ to be the map such that if X is a letter in the i th component of m and $|X|$ is in α_j , then $\theta(i)$ is equal to j . We define $\psi(p)$ to be (m, θ) .

We have the following lemma which corresponds to Lemma 5.7.

Lemma 8.2. *Let p and p' be elements of $\mathcal{P}(\alpha)$. If $p \sim p'$, then $\psi(p) \sim_M \psi(p')$.*

Proof. We omit the proof. It is similar to that of Lemma 5.7. $\square$

The following theorem is analogous to Theorem 5.8.

Theorem 8.3. *There is a bijection between the homotopy classes of $\mathcal{P}(\alpha)$ and the equivalence classes of $\mathcal{M}_R(\alpha)$ under $\sim_M$. In other words*

$$\mathcal{P}(\alpha)/\sim \cong \mathcal{M}_R(\alpha)/\sim_M.$$

We define $\mathcal{K}_M(\alpha)$ to be the set of equivalence classes of $\mathcal{M}_R(\alpha)$ under $\sim_K$. Then we can define the concept of i -reduction of an element of $\mathcal{K}_M(\alpha)$ as we did for $\mathcal{K}(\alpha)$. The proof that this concept is well-defined is almost identical to that of the case of $\mathcal{K}(\alpha)$ (Lemma 5.10).

It is easy to check that the Confluence condition proved in Lemma 5.12 also holds for $\mathcal{K}_M(\alpha)$. As the Finiteness condition also holds, we can use Newman's Diamond Lemma to get the following proposition corresponding to Proposition 5.14.

Proposition 8.4. *Each equivalence class of $\mathcal{K}_M(\alpha)$ under $\sim_M$ contains exactly one reduced element.*

For an element p of $\mathcal{P}$ we denote by $R_K(p)$ the reduced element in $\mathcal{K}_M(\alpha)$ corresponding to the element of $\mathcal{K}_M(\alpha)$ containing $\psi(p)$. The following theorem corresponds to Theorem 5.15.

Theorem 8.5. *For a nanophrase p in $\mathcal{P}(\alpha)$, $R_K(p)$ is a homotopy invariant of p .*

We extend the map s_i to elements of $\mathcal{M}_R(\alpha)$ in a natural way. For (m, θ) in $\mathcal{M}_R(\alpha)$ we define $s_i((m, \theta))$ to be the nanomultiphrase derived from (m, θ) by deleting all components for which the index of the component is mapped by θ to some integer other than i . By definition $s_i((m, \theta))$ contains the same number of phrases as (m, θ) .

Let (m, θ) be an element of $R_K(p)$. We write $\theta_R(p)$ for θ which is an invariant of p . We also define $P_{R,i}(p)$ to be $s_i((m, \theta))$. The nanomultiphrase $P_{R,i}(p)$ is a nanomultiphrase in $\mathcal{M}(\alpha_i)$ and is, modulo $\sim_i$, an invariant of p . Taken together, the map $\theta_R(p)$ and the set of nanomultiphrases $P_{R,i}(p)$ give a complete invariant for nanophrases.

Definitions of rank and homotopy rank naturally extend to nanomultiphrases and we use the same notation as we did for nanowords and nanophrases. Following the arguments about rank and homotopy rank in Section 5, it is easy to prove the following theorem which corresponds to Theorem 5.19.

Theorem 8.6. *Let (α, τ, S) be a composite homotopy data triple and let its prime factors be denoted by (α_i, τ_i, S_i) . Let p be a nanophrase over α . Then*

$$(8.1) \quad \text{hr}(p) = \sum_i \text{hr}(P_{R,i}(p)).$$

We finish by giving the following theorem.

Theorem 8.7. *Let (α, τ, S) be a composite homotopy data triple and let its prime factors be denoted by (α_i, τ_i, S_i) . Suppose, for each i , the homotopy given by (α_i, τ_i, S_i) is reduction decidable and equality decidable. Then the homotopy given by (α, τ, S) is also reduction decidable and equality decidable.*

Proof. The fact that (α, τ, S) is equality decidable is analogous to Theorem 6.2 and can be proved similarly.

Given a nanophrase p in $\mathcal{P}$, p is j -reducible if and only if all the components in the j th phrase of $\psi(p)$ are reducible. Since the homotopies given by (α_i, τ_i, S_i) are all reduction decidable, there is a finite time algorithm which determines the reducibility of all these components. Thus the homotopy given by (α, τ, S) is reduction decidable. $\square$

REFERENCES

- [1] P. M. Cohn, *Algebra, second edition, volume 3*, John Wiley & Sons, Chichester, 1997.
- [2] T. Fukunaga, *Homotopy classification of nanophrases with less than or equal to four letters*, arXiv:0904.3478.
- [3] ———, *On generalization of homotopy of words and its applications*, arXiv:0908.2899.
- [4] ———, *Homotopy classification of nanophrases in Turaev's theory of words*, J. Knot Theory Ramifications **18** (2009), no. 7, 901–915.
- [5] A. Gibson, *Homotopy invariants of Gauss phrases*, to appear in Indiana Univ. Math. J., arXiv:math.GT/0810.4389.
- [6] ———, *Homotopy invariants of Gauss words*, preprint, Tokyo Institute of Technology, 2008, arXiv:math.GT/0902.0062.
- [7] ———, *On tabulating virtual strings*, preprint, Tokyo Institute of Technology, 2008, arXiv:math.GT/0808.0064.

- [8] M. Goussarov, M. Polyak, and O. Viro, *Finite-type invariants of classical and virtual knots*, Topology **39** (September 2000), 1045–1068(24).
- [9] T. Kadokami, *Detecting non-triviality of virtual links*, J. Knot Theory Ramifications **12** (2003), no. 6, 781–803.
- [10] V. O. Manturov, *On free knots*, arXiv:math.GT/0901.2214.
- [11] ———, *On free knots and links*, arXiv:math.GT/0902.0127.
- [12] M. H. A. Newman, *On theories with a combinatorial definition of “equivalence”*, Ann. Math. **43** (1942), no. 2, 223–243.
- [13] V. Turaev, *Virtual strings*, Ann. Inst. Fourier (Grenoble) **54** (2004), no. 7, 2455–2525.
- [14] ———, *Knots and words*, Int. Math. Res. Not. (2006), Art. ID 84098, 23.
- [15] ———, *Topology of words*, Proc. Lond. Math. Soc. (3) **95** (2007), no. 2, 360–412.

DEPARTMENT OF MATHEMATICS, TOKYO INSTITUTE OF TECHNOLOGY, OH-OKAYAMA, MEGURO,
TOKYO 152-8551, JAPAN

E-mail address: `gibson@math.titech.ac.jp`